



Need more money for college expenses?

The CLC Private LoanSM can get you up to
\$40,000 a year for college-related expenses.

Here's why the CLC Private LoanSM is a smart choice:

- ✓ Approved borrowers are sent a check within four business days
- ✓ Get \$1000 - \$40,000 each year
- ✓ Interest rates as low as prime + 0% (8.66% APR)
- ✓ Quick and easy approval process
- ✓ No payments until after graduating or leaving school

CLICK HERE

or call **800.311.9615.**

*We are available 24 hours
a day, 7 days a week.*

4

Rational Expressions

Doug Nance opens the valve to fill his new swimming pool. Without his knowledge, the drain to empty the pool has been left open. If the pool would normally fill in 10 hours and the drain can empty the full pool in 15 hours, how long does it take to fill the pool?



4-1 ■ Fundamental principle of rational expressions

The rational expression

In chapter 1, we defined a rational number to be any number that can be expressed as a quotient of two integers where the denominator is not zero. That is, the set of rational numbers are the elements in the set

$$\left\{ \frac{p}{q} \mid p \text{ and } q \text{ are integers, } q \neq 0 \right\}$$

If this definition is extended to polynomials, the result is called a **rational expression**.

Definition

A **rational expression** is any algebraic expression that can be written as a quotient of two polynomials, where the denominator is not equal to zero.

That is, rational expressions are the elements in the set

$$\left\{ \frac{P}{Q} \mid P \text{ and } Q \text{ are polynomials, } Q \neq 0 \right\}$$

The following are rational expressions:

$$\frac{2x}{3}, \quad \frac{2x-3}{x^2+6x-1}, \quad \frac{5x+1}{3}, \quad \text{and} \quad 5x^3-3x+1$$

Note $5x^3 - 3x + 1$ is a rational expression since it can be written

$$\frac{5x^3 - 3x + 1}{1}$$

Recall that division by zero is not defined. The rational expression becomes meaningless for those values of the variable for which the denominator is zero. For example, the rational expression

$$\frac{3x - 2}{x - 6}$$

is meaningless if $x = 6$, because then $x - 6 = 6 - 6 = 0$. Thus we say that

$$\frac{3x - 2}{x - 6}, x \neq 6$$

is a rational expression.

Note The restrictions on the variable are found by setting all factors of the denominator that contain the variable equal to zero and solving the resulting equation(s) for the variable.

Domain of a rational expression

We call the set of all those values that the variable in a rational expression can assume the **domain** of the rational expression.

Definition of the domain

The **domain** of a rational expression is the set of all replacement values of the variable for which the expression is defined.

Thus, given the rational expression

$$\frac{3x - 2}{x - 6}$$

the domain is defined to be the set of all real numbers except 6. In set-builder notation, we symbolize the domain as follows:

$$\text{domain} = \{x | x \in R, x \neq 6\}$$

When the factored denominator consists of more than one factor containing the variable, we use the following property to determine restrictions on the variable.

Zero product property

If P and Q are polynomials, then

$$P \cdot Q = 0$$

if and only if $P = 0$ or $Q = 0$.

Concept

The product of two polynomials is zero provided one or both of the polynomials is equal to zero.

Example 4-1 A

Determine the domain of the following rational expressions. Express this in set-builder notation.

1. $\frac{x-5}{x+7}$

$$\begin{aligned}x+7 &= 0 \\x &= -7\end{aligned}$$

Set the denominator equal to 0
Solve the equation

The number -7 cannot be used as a replacement for x .

$$\text{Domain} = \{x | x \in R, x \neq -7\}$$

2. $\frac{4x-3}{x^2-16}$

$$\begin{aligned}x^2 - 16 &= (x+4)(x-4) \\x+4 &= 0 \quad \text{or} \quad x-4 = 0 \\x &= -4 \quad \quad \quad x = 4\end{aligned}$$

Factor the denominator
Set each factor equal to 0

The numbers -4 and 4 cannot be used as replacements for x .

$$\text{Domain} = \{x | x \in R, x \neq -4, x \neq 4\}$$

3. $\frac{5y^2}{y^2+9}$

Since $y^2 + 9$ is positive for every value of y , then $y^2 + 9$ is *never* equal to zero. Hence, the domain $= \{y | y \in R\}$.

► **Quick check** Determine the domain of $\frac{3y-2}{y^2-1}$ and express the domain in set-builder notation.

Reducing to lowest terms

The methods used to simplify rational expressions by reducing them to lowest terms are similar to those used to simplify fractions. A rational expression is reduced to lowest terms when there is no polynomial factor (other than 1 or -1) that is common to both the numerator and the denominator. This is accomplished by factoring the numerator and the denominator and removing a factor of 1. Recall

$$\frac{8}{8} = 1; \quad \frac{-9}{-9} = 1; \quad \frac{x-2}{x-2} = 1, \text{ if } x \neq 2$$

Given polynomials P , Q , and R ,

$$\frac{PR}{QR} = \frac{P \cdot R}{Q \cdot R} = \frac{P}{Q} \cdot \frac{R}{R} = \frac{P}{Q} \cdot 1 = \frac{P}{Q}$$

Thus,

$$\frac{(x+2)(x-3)}{(2x-1)(x-3)} = \frac{x+2}{2x-1} \cdot \frac{x-3}{x-3} = \frac{x+2}{2x-1} \cdot 1 = \frac{x+2}{2x-1} \quad \left(x \neq \frac{1}{2} \text{ or } 3\right)$$

We have applied the fundamental principle of rational expressions which is now stated.

Fundamental principle of rational expressions

If P , Q , and R are polynomials, $Q \neq 0$ and $R \neq 0$, then

$$\frac{P}{Q} = \frac{P \cdot R}{Q \cdot R} \quad \text{and} \quad \frac{P \cdot R}{Q \cdot R} = \frac{P}{Q}$$

Concept

Both the numerator and the denominator of a rational expression may be multiplied or divided by the same nonzero polynomial without changing the value of the expression.

By this principle, the rational expression

$$\begin{aligned} \frac{x-3}{x^2-9} &= \frac{x-3}{(x+3)(x-3)} && \text{Factor the denominator} \\ &= \frac{1}{x+3} \quad (x \neq -3, x \neq 3) && \text{Divide the numerator and the denominator by } x-3 \end{aligned}$$

Note The domain of the expression excludes -3 and 3 even though the factor $x-3$ is not in the resulting expression.

We use the **fundamental principle of rational expressions** to simplify rational expressions by reducing them to lowest terms. This is accomplished by dividing the numerator and the denominator by all common factors.

To reduce a rational expression to lowest terms

1. Completely factor the numerator and the denominator.
2. Divide the numerator and the denominator by the greatest common factor.

Example 4-1 B

Simplify each rational expression by reducing it to lowest terms. State any necessary restrictions on the variables.

$$\begin{aligned} 1. \quad \frac{16x^2}{24x^4} &= \frac{2(8x^2)}{3x^2(8x^2)} && \begin{array}{l} \text{Factor the numerator and the denominator} \\ \text{Greatest common factor} \end{array} \\ &= \frac{2}{3x^2} \quad (x \neq 0) && \begin{array}{l} \text{Divide the numerator and the denominator} \\ \text{by } 8x^2 \end{array} \\ 2. \quad \frac{6x-18}{5x-15} &= \frac{6(x-3)}{5(x-3)} && \text{Factor the numerator and the denominator} \\ &= \frac{6}{5} \quad (x \neq 3) && \text{Divide the numerator and the denominator} \\ &&& \text{by the common factor } x-3 \end{aligned}$$

$$3. \frac{y-4}{4-y}$$

We note that the numerator and the denominator are exactly the same except for the signs. That is, $4 - y = -y + 4 = (-1)(y - 4)$. The given expression can be written in lowest terms by multiplying the numerator and the denominator by -1 .

$$\begin{aligned} &= \frac{(-1)(y-4)}{(-1)(4-y)} \\ &= \frac{(-1)(y-4)}{y-4} \\ &= \frac{-1}{1} \\ &= -1 \quad (y \neq 4) \end{aligned}$$

Multiply numerator and denominator by -1

Multiply in the denominator, $(-1)(4 - y) = y - 4$

Reduce by the common factor $y - 4$

$$4. \frac{x^2 - x - 12}{20 - x - x^2} = \frac{(x-4)(x+3)}{(4-x)(5+x)}$$

Factor the numerator and the denominator

We note opposite factors $x - 4$ and $4 - x$ in the numerator and the denominator.

$$\begin{aligned} &= \frac{(-1)(x-4)(x+3)}{(-1)(4-x)(5+x)} \\ &= \frac{(-1)(x-4)(x+3)}{(x-4)(5+x)} \\ &= \frac{(-1)(x+3)}{1(5+x)} \\ &= \frac{x+3}{x+5} \quad (x \neq 4, x \neq -5) \end{aligned}$$

Multiply numerator and denominator by -1

$(-1)(4 - x) = x - 4$

Reduce by $x - 4$

$-\frac{a}{b} = \frac{a}{b}$ and $5 + x = x + 5$

► **Quick check** Simplify $\frac{x^2 + 3x - 28}{x^2 - 16}$ by reducing to lowest terms. State any necessary restrictions on the variables.

Mastery points

Can you

- Determine the domain of a rational expression?
- Apply the fundamental principle of rational expressions to reduce a rational expression to lowest terms?

Exercise 4-1

State the domain of the given rational expression in set-builder notation. See example 4-1 A.

Example $\frac{3y-2}{y^2-1}$

Solution
$$= \frac{3y - 2}{(y + 1)(y - 1)}$$

Factor the denominator

$$y + 1 = 0 \quad \text{or} \quad y - 1 = 0$$

$$y = -1 \quad \quad \quad y = 1$$

Set each factor in the denominator equal to 0

Solve each equation

The numbers 1 and -1 cannot be used as replacements for y .

$$\text{Domain} = \{y | y \in R, y \neq -1, y \neq 1\}$$

1. $\frac{3}{x-4}$

2. $\frac{5}{x-8}$

3. $\frac{x}{x+1}$

4. $\frac{5y}{y+9}$

5. $\frac{3z}{2z-3}$

6. $\frac{5x}{8x-5}$

7. $\frac{x-3}{7x+4}$

8. $\frac{p+9}{6p+5}$

9. $\frac{2x-3}{x^2-4x}$

10. $\frac{m+5}{3m^2+6m}$

11. $\frac{4x-7}{x^2+8x+16}$

12. $\frac{7-5x}{x^2-14x+49}$

13. $\frac{2y-5}{4y^2-25}$

14. $\frac{3x+4}{9x^2-16}$

15. $\frac{8x^2+1}{2x^2-5x-3}$

16. $\frac{p^2-7p+1}{4p^2+p-3}$

17. $\frac{4x^2-2x+1}{x^2+4}$

18. $\frac{x-3}{x^2+5}$

Simplify the given rational expression by reducing to lowest terms. State any necessary restrictions on the variable. See example 4-1 B.

Example
$$\frac{x^2+3x-28}{x^2-16}$$

Solution
$$= \frac{(x-4)(x+7)}{(x-4)(x+4)}$$

Greatest common factor

Factor the numerator and the denominator

$$= \frac{x+7}{x+4} \quad (x \neq -4)$$

Greatest common factor

Reduce by $x-4$

19. $\frac{25x^3}{15x^4}$

20. $\frac{36y^4}{-42y^2}$

21. $\frac{p^3q^5}{pq^7}$

22. $\frac{-m^5n^3}{-m^8n}$

23. $\frac{a^2b^4c^3}{-abc^7}$

24. $\frac{-x^4y^6z}{x^3y^5z}$

25. $-\frac{27mn^3p^7}{36m^4np^5}$

26. $\frac{72p^3qr^4}{-63pqr}$

27. $\frac{3x-6}{7x-14}$

28. $\frac{8a+12}{6a+9}$

29. $\frac{15a}{5a^2-10a}$

30. $\frac{-36x}{42x^3+24x}$

31. $\frac{12x^4-12x^3}{6x^3}$

32. $\frac{24x^2-36x}{32x^2}$

33. $\frac{8y^2-8}{6y-6}$

34. $\frac{6x+6}{3x-3}$

35. $\frac{5x-5y}{y-x}$

36. $\frac{9b-9a}{a-b}$

37. $\frac{a^2-9}{4a+12}$

38. $\frac{6a+3b}{4a^2-b^2}$

39. $\frac{4y^2-1}{1+2y}$

40. $\frac{64-49p^2}{7p-8}$

41. $\frac{x^3+8}{x+2}$

42. $\frac{a^3-64}{a-4}$

43. $\frac{3x-3y}{2y^3-2x^3}$

44. $\frac{5a^3+5b^3}{7b+7a}$

45. $\frac{y^2-49}{y^2+14y+49}$

46. $\frac{a^2-10a+25}{a^2-25}$

47. $\frac{m^2-4m-12}{m^2-m-6}$

48. $\frac{n^2-4n+3}{n^2-5n+6}$

49. $\frac{y^2-4y-32}{y^2-y-20}$

50. $\frac{x^2-x-42}{x^2+12x+36}$

51. $\frac{2a^2-3a+1}{2a^2+a-1}$

52. $\frac{3b^2-10b+3}{3b^2-7b+2}$

53. $\frac{2x^2+3x-9}{4x^2+13x+3}$

54. $\frac{8y^2-22y+5}{4y^2-15y-4}$

55. $\frac{12m^2+17m+6}{8m^2+10m+3}$

Love The Taste. Taste The Love.

At Culver's® we can't think of anything better than serving up our creamy frozen custard and delicious classics cooked fresh the minute you order them. Which is why when we bring them to your table, they're always accompanied by a warm smile and a friendly offer to see if there's anything else we can get for you. So come on into your neighborhood Culver's and see for yourself. You might just be in love by the time you leave.



56. $\frac{6x^2 + 17x + 7}{12x^2 + 13x - 35}$

57. $\frac{4a^2 + 2a - 12}{6a^2 - 7a - 3}$

58. $\frac{6x^2 - x - 1}{4x^2 + 18x - 10}$

59. $\frac{5y^2 - 10y - 15}{4y^2 - 36}$

60. $\frac{7x^2 - 7}{3x^2 - 15x + 12}$

61. $\frac{3a^2 + 16a - 12}{6 - 7a - 3a^2}$

62. $\frac{a^2 + 2ab - 24b^2}{a^2 - 8ab + 16b^2}$

63. $\frac{p^2 + 7pq + 12q^2}{p^2 + 5pq + 6q^2}$

Review exercises

Perform the indicated operations. See section 1-4.

1. $\frac{3}{4} - \frac{5}{6}$

2. $\frac{5}{6} \div \frac{10}{12}$

3. $\frac{7}{8} + \frac{3}{4}$

Find the solution set of the following equations. See section 2-1.

4. $4(x - 2) + 3 = 2x - 1$

5. $\frac{3}{2}x - 2 = \frac{5}{6}x + 1$

6. Solve $4x + 2a = 6x + 5$ for x . See section 2-2.

4-2 ■ Multiplication and division of rational expressions**Multiplication**

Recall that the product of two rational numbers is obtained by multiplying the numerators and placing this product over the product of the denominators. This process is indicated by the **multiplication property of rational numbers** stated here.

Multiplication property of rational numbers

If a , b , c , and d are real numbers, $b \neq 0$ and $d \neq 0$, then

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{a \cdot c}{b \cdot d}$$

To illustrate,

$$\frac{3}{4} \cdot \frac{5}{7} = \frac{3 \cdot 5}{4 \cdot 7} = \frac{15}{28}$$

To multiply two or more rational expressions we use the **multiplication property of rational expressions**.

Multiplication property of rational expressions

If P , Q , R , and S are polynomials, $Q \neq 0$ and $S \neq 0$, then

$$\frac{P}{Q} \cdot \frac{R}{S} = \frac{P \cdot R}{Q \cdot S}$$

Concept

When multiplying two rational expressions, multiply the two numerators to determine the numerator of the product and multiply the two denominators to determine the denominator of the product.

For example,

$$\frac{x-3}{x+1} \cdot \frac{x+2}{x-1} = \frac{(x-3)(x+2)}{(x+1)(x-1)} = \frac{x^2 - x - 6}{x^2 - 1}$$

Since we want the resulting product in its simplest form, any possible reduction by common factors should be performed **before** the multiplication of the numerators and the denominators takes place.

To multiply rational expressions

1. Completely factor the numerators and the denominators where possible.
2. Write the factors in the numerators and the denominators as a single rational expression.
3. Divide the numerator and the denominator by all common factors.
4. Multiply the remaining factors in the numerator and in the denominator to obtain a rational expression in lowest terms.

Example 4-2 A

Find the indicated products reduced to lowest terms. Place restrictions on the variables.

$$1. \frac{6x^2}{15} \cdot \frac{10}{3x^3} = \frac{6x^2 \cdot 10}{15 \cdot 3x^3}$$

$$= \frac{4 \cdot \cancel{15x^2}}{3x \cdot \cancel{15x^2}}$$

Greatest common factor
Factor numerator and denominator
Greatest common factor

$$= \frac{4}{3x} \quad (x \neq 0)$$

Divide numerator and denominator by the common factor $15x^2$

$$2. \frac{3x^2 - 6x}{x + 5} \cdot \frac{x^2 - 25}{x^2 - 10x + 25}$$

$$= \frac{3x(x-2)}{x+5} \cdot \frac{(x+5)(x-5)}{(x-5)(x-5)}$$

Factor where possible

$$= \frac{3x(x-2)(x+5)(x-5)}{(x+5)(x-5)(x-5)}$$

Multiply numerators and denominators

$$= \frac{3x(x-2)}{x-5}$$

Divide numerator and denominator by the common factors $x+5$ and $x-5$

$$= \frac{3x^2 - 6x}{x-5} \quad (x \neq -5, 5)$$

Multiply in the numerator

Note You may leave your answer in factored form $\frac{3x(x-2)}{x-5}$ or multiply as indicated to obtain $\frac{3x^2 - 6x}{x-5}$; we will show both answers.

$$\begin{aligned}
 3. \quad & \frac{3x-1}{x-4} \cdot \frac{16-x^2}{3x^2+14x-5} \\
 &= \frac{3x-1}{x-4} \cdot \frac{(4-x)(4+x)}{(3x-1)(x+5)} \\
 &= \frac{(3x-1)(4-x)(4+x)}{(x-4)(3x-1)(x+5)}
 \end{aligned}$$

Since there are opposite factors $4-x$ and $x-4$, we multiply the numerator and the denominator by -1 .

$$\begin{aligned}
 &= \frac{(-1)(4-x)(3x-1)(4+x)}{(-1)(x-4)(3x-1)(x+5)} \\
 &= \frac{(-1)(4-x)(3x-1)(4+x)}{(4-x)(3x-1)(x+5)} && (-1)(x-4) = 4-x \\
 &= \frac{(-1)(4+x)}{x+5} && \text{Reduce by the common factors } 4-x \text{ and } 3x-1 \\
 &= \frac{-x-4}{x+5} \quad \left(x \neq 4, \frac{1}{3}, -5 \right)
 \end{aligned}$$

► **Quick check** Find the product of $\frac{4x^2-12x}{x+5} \cdot \frac{x^2-25}{x^2-6x+9}$ and reduce to lowest terms. Place restrictions on the variable.

Division of rational expressions

Recall that the reciprocal of any nonzero real number b is the real number $\frac{1}{b}$ found by “inverting” the number. In like fashion, we invert rational expressions to obtain their reciprocals. That is,

$$\frac{x-1}{x+3} \quad \text{and} \quad \frac{x+3}{x-1} \quad \text{are reciprocals.}$$

We use reciprocals in the division of rational expressions as we did when dividing rational numbers.

Division property of rational numbers

If a , b , c , and d are real numbers and $b \neq 0$, $c \neq 0$, and $d \neq 0$, then

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} = \frac{a \cdot d}{b \cdot c}$$

To illustrate,

$$\frac{5}{8} \div \frac{3}{7} = \frac{5}{8} \cdot \frac{7}{3} = \frac{5 \cdot 7}{8 \cdot 3} = \frac{35}{24}$$

We can extend this property to rational expressions.

Division property of rational expressions

If P , Q , R , and S are polynomials, $Q \neq 0$, $R \neq 0$, and $S \neq 0$, then

$$\frac{P}{Q} \div \frac{R}{S} = \frac{P}{Q} \cdot \frac{S}{R} = \frac{P \cdot S}{Q \cdot R}$$

Concept

When dividing two rational expressions, the quotient is obtained by multiplying the first expression by the reciprocal of the second expression.

Example 4-2 B

Find each of the indicated quotients reduced to lowest terms. Assume no denominator equals zero.

$$\begin{aligned}
 1. \quad \frac{36p^3}{7q} \div \frac{15p}{14q^2} &= \frac{36p^3}{7q} \cdot \frac{14q^2}{15p} && \text{Multiply by the reciprocal of } \frac{15p}{14q^2} \\
 &= \frac{36 \cdot 14 \cdot p^3 q^2}{7 \cdot 15 \cdot pq} && \text{Multiply numerators and denominators} \\
 &= \frac{3 \cdot 12 \cdot 7 \cdot 2 \cdot p^3 q^2}{7 \cdot 3 \cdot 5 \cdot pq} && \text{Factor numerator and denominator} \\
 &= \frac{24p^2 q \cdot (21pq)}{5 \cdot (21pq)} && \text{Greatest common factor} \\
 &= \frac{24p^2 q}{5} && \text{Greatest common factor} \\
 &&& \text{Reduce by } 21pq
 \end{aligned}$$

$$\begin{aligned}
 2. \quad \frac{2x+1}{x^2-9} \div \frac{2x^2+7x+3}{x-3} &= \frac{2x+1}{x^2-9} \cdot \frac{x-3}{2x^2+7x+3} && \text{Multiply by the reciprocal of } \frac{2x^2+7x+3}{x-3} \\
 &= \frac{(2x+1)(x-3)}{(x+3)(x-3)(2x+1)(x+3)} && \text{Factor and multiply numerators and denominators} \\
 &= \frac{1}{(x+3)(x+3)} && \text{Reduce by } (x-3)(2x+1) \\
 &= \frac{1}{x^2+6x+9} && \text{Multiply as indicated}
 \end{aligned}$$

$$\begin{aligned}
 3. \quad \frac{x^2+2x}{5x} \div (2x^2+x-6) &= \frac{x^2+2x}{5x} \div \frac{2x^2+x-6}{1} && \text{Since } 2x^2+x-6 = \frac{2x^2+x-6}{1}, \text{ then} \\
 \frac{x^2+2x}{5x} \div \frac{2x^2+x-6}{1} &= \frac{x^2+2x}{5x} \cdot \frac{1}{2x^2+x-6} && \text{Multiply by the reciprocal of } \frac{2x^2+x-6}{1} \\
 &= \frac{x(x+2)}{5x(2x-3)(x+2)} && \text{Factor and multiply numerators and denominators} \\
 &= \frac{1}{5(2x-3)} && \text{Reduce by } x(x+2) \\
 &= \frac{1}{10x-15} && \text{Multiply as indicated}
 \end{aligned}$$

► **Quick check** Find the quotient of $\frac{4x+3}{x^2-49} \div \frac{4x^2-13x-12}{x+7}$ and reduce to lowest terms. Assume no denominator equals zero.

Mastery points

Can you

- Multiply rational expressions?
- Divide rational expressions?

Exercise 4-2

Find each indicated product in lowest terms. Place restrictions on the variable. See example 4-2 A.

Example $\frac{4x^2-12x}{x+5} \cdot \frac{x^2-25}{x^2-6x+9}$

Solution

$$\begin{aligned} &= \frac{4x(x-3)}{x+5} \cdot \frac{(x+5)(x-5)}{(x-3)(x-3)} \\ &= \frac{4x(x-3)(x+5)(x-5)}{(x+5)(x-3)(x-3)} \\ &= \frac{4x(x-5)\cancel{(x-3)}\cancel{(x+5)}}{(x-3)\cancel{(x-3)}\cancel{(x+5)}} \\ &= \frac{4x(x-5)}{x-3} \\ &= \frac{4x^2-20x}{x-3} \quad (x \neq -5, 3) \end{aligned}$$

Factor the numerator and the denominator

Multiply the numerators and the denominators

Greatest common factors

Commute

Greatest common factors

Reduce by $(x-3)(x+5)$

Multiply in the numerator

1. $\frac{8}{9x^2} \cdot \frac{12x}{4}$

2. $\frac{12b}{7} \cdot \frac{28}{4b^3}$

3. $\frac{5x^2}{9y^3} \cdot \frac{18y}{20x^4}$

4. $\frac{15n}{28m} \cdot \frac{14m^3}{30n^2}$

5. $\frac{16c}{42ab^2} \cdot \frac{3a^3b}{8c^4}$

6. $\frac{12xy}{25z^3} \cdot \frac{5z}{18x^3y^2}$

7. $16a^2b^3 \cdot \frac{3}{24ab}$

8. $8x^3yz^2 \cdot \frac{15}{36xyz}$

9. $\frac{4a+8}{3a-12} \cdot \frac{a-2}{5a+10}$

10. $\frac{y+3}{8y-16} \cdot \frac{3y-6}{4y+12}$

11. $(a+3) \cdot \frac{a-6}{6a+18}$

12. $\frac{x-3}{5x-25} \cdot (5-x)$

13. $\frac{p^2+2p+1}{1-4p} \cdot \frac{16p^2-1}{p^2-1}$

14. $\frac{m^2-9}{3m+4} \cdot \frac{9m^2-16}{m^2+6m+9}$

15. $\frac{x^2-9x+20}{x^2-5x+6} \cdot \frac{x^2-3x+2}{x^2-5x+4}$

16. $\frac{y^2+3y-4}{y^2+2y-3} \cdot \frac{y^2-y-6}{y^2+y-12}$

17. $\frac{2a^2-a-6}{3a^2-4a+1} \cdot \frac{3a^2+7a+2}{2a^2+7a+6}$

18. $\frac{4b^2+8b+3}{2b^2-5b-12} \cdot \frac{b^2-16}{2b^2+7b+3}$

19. $\frac{x^3+8}{x^2-4} \cdot \frac{x^2-4x+4}{2x^2+3x-14}$

20. $\frac{4x^2-49}{8x^3+27} \cdot \frac{4x^2+12x+9}{2x^2-13x+21}$

21. $\frac{y-x}{x^2+3xy+2y^2} \cdot \frac{x^2+2xy+y^2}{x^2-y^2}$

22. $\frac{b^2-a^2}{3a^2+ab-2b^2} \cdot \frac{6a^2-ab-2b^2}{a^2-2ab+b^2}$

Find each of the indicated quotients in lowest terms. Assume all denominators are nonzero. See example 4-2 B.

Example $\frac{4x+3}{x^2-49} \div \frac{4x^2-13x-12}{x+7}$

Solution
$$\begin{aligned} &= \frac{4x+3}{x^2-49} \cdot \frac{x+7}{4x^2-13x-12} \\ &= \frac{(4x+3)(x+7)}{(x+7)(x-7)(4x+3)(x-4)} \\ &= \frac{(4x+3)(x+7)}{(4x+3)(x+7)(x-7)(x-4)} \\ &= \frac{1}{(x-7)(x-4)} \\ &= \frac{1}{x^2-11x+28} \end{aligned}$$

Multiply by the reciprocal

Factor the denominator and multiply

Commute

Reduce by GCF $(4x+3)(x+7)$

Multiply in the denominator

23. $\frac{9a^2b}{8ab^3} \div \frac{18ab^3}{16a^2b^2}$

24. $\frac{7ab^3}{10x^2y^3} \div \frac{21a^2b^2}{15x^3y^2}$

25. $\frac{r+4}{r^2-1} \div \frac{r^2-16}{r+1}$

26. $\frac{4x+4y}{x-3} \div \frac{x+y}{x^2-9}$

27. $\frac{6-2x}{2x+8} \div (9-3x)$

28. $\frac{x^2-25}{2x+10} \div (x^2-10x+25)$

29. $(x^2-2x-3) \div \frac{4x-12}{x^2-1}$

30. $(4x^2-9) \div \frac{4x+6}{x+3}$

31. $\frac{x^3-125}{2x+1} \div \frac{x^2-25}{2x^2-9x-5}$

32. $\frac{3x^2+10x-8}{5x-15} \div \frac{x^3+64}{x^2-9}$

33. $\frac{56-x-x^2}{x^2-6x-7} \div \frac{x^2+4x-21}{x^2-4x-5}$

34. $\frac{a^2+16a+63}{20-a-a^2} \div \frac{a^2-5a-84}{a^2-a-12}$

35. $\frac{2x^2+5x-12}{3x^2-8x-16} \div \frac{2x^2+3x-9}{3x^2+13x+12}$

36. $\frac{6x^2-5x-4}{8x^2+10x+3} \div \frac{3x^2+14x-24}{4x^2-12x-7}$

Find each indicated product or quotient in lowest terms. Assume all denominators are nonzero. See examples 4-2 A and B.

37. $\frac{a^2-b^2}{2a+4b} \cdot \frac{a+2b}{a+b}$

39. $\frac{n^2-m^2}{2m-3n} \div \frac{m-n}{4m^2-9n^2}$

41. $\frac{a^2-8a+15}{a^2+7a+6} \cdot \frac{a^2-6a-7}{a^2-9} \div \frac{a^2-12a+35}{a^2-36}$

43. $\frac{2x^2+5x-3}{3x^2-10x-8} \div \frac{2x^2-7x+3}{3x^2-x-2} \cdot \frac{x^2-6x+9}{x^2-2x+1}$

45. $\frac{27y^3-8x^3}{x+y} \div \frac{2x-3y}{x^3+y^3}$

38. $\frac{4x+4y}{x-3y} \cdot \frac{x^2-9y^2}{x+y}$

40. $\frac{p^2-q^2}{q-3p} \div \frac{p^2+2pq+q^2}{9p^2-q^2}$

42. $\frac{x^2+3x-28}{x^2-8x+7} \div \frac{x^2-7x+12}{x^2-x-42} \cdot \frac{2x-8}{x^2-36}$

44. $\frac{2x^2+x-6}{5x^2+7x-6} \cdot \frac{20x^2-7x-3}{6x^2-25x+4} \div \frac{4x^2-11x-3}{6x^2-19x+3}$

46. $\frac{a^3+64b^3}{16b^2-a^2} \cdot \frac{a+4b}{a^2+3ab-4b^2}$



Extra Credit Rocks

Sign up for a Discover® Student Card today and enjoy:

- 0% Intro APR* on Purchases for 6 Months
- No Annual Fee
- Easiest Online Account Management Options
- Full 5% *Cashback Bonus*®* on Get More purchases in popular categories all year
- Up to 1% *Cashback Bonus*®* on all your other purchases
- Unlimited cash rewards that never expire as long as you use your Card

APPLY NOW

DISCOVER
CARD

*View Discover® Card Rates, Fees, Rewards and Other Important Information.

$$47. \frac{ab - a + 3b - 3}{ab + a - 4b - 4} \div \frac{ab - 2a + 3b - 6}{ab + 2a - 4b - 8}$$

$$49. \frac{xz - xw + yz - yw}{xz + xw - yz - yw} \div \frac{xz - xw - yz + yw}{xz + xw + yz + yw}$$

$$48. \frac{mn - n + 4m - 4}{mn - 3n + 4m - 12} \div \frac{mn - 3n + 2m - 6}{mn + 2m + n + 2}$$

$$50. \frac{pr + ps + qr + qs}{pr - ps - qr + qs} \div \frac{xr + xs - yr - ys}{xr - xs + yr - ys}$$

Review exercises

Add or subtract the following fractions. See section 1-4.

$$1. \frac{11}{12} + \frac{7}{8}$$

$$2. \frac{14}{15} - \frac{3}{5}$$

Find the solution set of the following inequalities. See section 2-5.

$$3. 5y - 2 \leq y + 6$$

$$4. -7 \leq 2y - 1 < 1$$

Find the following products. See section 3-2.

$$5. 4x^3(2x^2 - x + 6)$$

$$6. (4y + 1)(y - 4)$$

4-3 ■ Addition and subtraction of rational expressions

When we add or subtract a pair of rational numbers that have the same denominator, such as $\frac{3}{4} + \frac{5}{4}$, we use the property of adding or subtracting fractions.

Property of adding or subtracting rational numbers

If a , b , and c are real numbers, $b \neq 0$, then

$$\frac{a}{b} + \frac{c}{b} = \frac{a+c}{b} \quad \text{and} \quad \frac{a}{b} - \frac{c}{b} = \frac{a-c}{b}$$

For example,

$$\frac{3}{4} + \frac{5}{4} = \frac{3+5}{4} = \frac{8}{4} = 2 \quad \text{and} \quad \frac{7}{8} - \frac{1}{8} = \frac{7-1}{8} = \frac{6}{8} = \frac{3}{4}$$

We can extend this property to rational expressions.

Property of adding or subtracting rational expressions

If P , Q , and R are polynomials, $Q \neq 0$, then

$$\frac{P}{Q} + \frac{R}{Q} = \frac{P+R}{Q} \quad \text{and} \quad \frac{P}{Q} - \frac{R}{Q} = \frac{P-R}{Q}$$

Concept

To add, or subtract, two rational expressions having the same denominators, add, or subtract, the numerators and place this sum or difference over the common denominator.

Example 4-3 A

Find each indicated sum or difference in lowest terms. Assume all denominators are nonzero.

$$1. \frac{3x+2}{x-2} + \frac{4x-5}{x-2}$$

$$= \frac{(3x+2) + (4x-5)}{x-2}$$

Add the numerators.

$$= \frac{3x+2+4x-5}{x-2}$$

Remove parentheses.

$$= \frac{7x-3}{x-2}$$

Combine like terms in the numerator.

$$2. \frac{3x-2y}{x^2-y^2} - \frac{2x-3y}{x^2-y^2}$$

$$= \frac{(3x-2y) - (2x-3y)}{x^2-y^2}$$

Subtract the numerators.

$$= \frac{3x-2y-2x+3y}{x^2-y^2}$$

Remove parentheses.

$$= \frac{x+y}{x^2-y^2}$$

Combine like terms.

$$= \frac{x+y}{(x+y)(x-y)}$$

Factor the denominator.

$$= \frac{1}{x-y}$$

Reduce by $(x+y)$ (always reduce when possible).

Note When we write the sum or the difference of the numerators, it is important that we write each numerator in parentheses to avoid making a common mistake when we subtract. Failure to do this would have caused the numerator in example 2 to become $3x-2y-2x-3y = x-5y$.

► **Quick check** Find the difference of $\frac{4x-3}{x^2-4} - \frac{3x-1}{x^2-4}$ in lowest terms.

Assume the denominator is nonzero.

When adding, or subtracting, two rational expressions $\frac{P}{Q}$ and $\frac{R}{S}$ that *do not* have the same denominator, we can add or subtract them in the following way:

Property of adding or subtracting rational expressions with different denominators

If P , Q , R , and S are polynomials, $Q \neq 0$ and $S \neq 0$, then

$$\frac{P}{Q} + \frac{R}{S} = \frac{P \cdot S + Q \cdot R}{Q \cdot S} \quad \text{and} \quad \frac{P}{Q} - \frac{R}{S} = \frac{P \cdot S - Q \cdot R}{Q \cdot S}$$

The following examples illustrate the use of this property for rational expressions.

$$\begin{aligned}
 1. \quad \frac{x+1}{x-2} + \frac{x-3}{x+4} &= \frac{(x+1)(x+4) + (x-2)(x-3)}{(x-2)(x+4)} && \text{Multiply as indicated} \\
 &= \frac{(x^2 + 5x + 4) + (x^2 - 5x + 6)}{(x-2)(x+4)} && \text{Combine like terms} \\
 &= \frac{2x^2 + 10}{(x-2)(x+4)} \quad (x \neq 2, x \neq -4)
 \end{aligned}$$

$$\begin{aligned}
 2. \quad \frac{x}{x-7} - \frac{2x+1}{x-9} &= \frac{x \cdot (x-9) - (x-7)(2x+1)}{(x-7)(x-9)} && \text{Multiply as indicated} \\
 &= \frac{(x^2 - 9x) - (2x^2 - 13x - 7)}{(x-7)(x-9)} && \text{Remove parentheses and change signs} \\
 &= \frac{x^2 - 9x - 2x^2 + 13x + 7}{(x-7)(x-9)} \\
 &= \frac{-x^2 + 4x + 7}{(x-7)(x-9)} \quad (x \neq 7, x \neq 9) && \text{Combine like terms}
 \end{aligned}$$

Note We do not multiply in the denominator to aid in recognizing common factors for reducing to lowest terms. Always check the numerator for common factors.

This method is useful when adding or subtracting only two rational expressions and when the denominators have no common factors. When rational numbers or rational expressions do not have the same denominator, we apply the fundamental principle of rational expressions to obtain equivalent expressions (expressions that name the same number) with a common denominator. To do this, we must find the *least common multiple* (LCM) of the denominators.

Least common multiple

The least common multiple (LCM) of a set of expressions is the least expression that is exactly divisible by the expressions.

We use the following procedure to find the LCM of a set of polynomials.

Finding the LCM of a set of polynomials

1. Completely factor the polynomials.
2. The LCM consists of the product of all distinct factors of the polynomials, each raised to the greatest power to which it appears in any one of the factorizations.

Example 4-3 B

Find the least common multiple.

1. $24x^3$, $15x^2$, $10x$

$$\left. \begin{aligned} 24x^3 &= 2^3 \cdot 3 \cdot x^3 \\ 15x^2 &= 3 \cdot 5 \cdot x^2 \\ 10x &= 2 \cdot 5 \cdot x \end{aligned} \right\} \text{Completely factor each polynomial}$$

Choose the greatest power of each different factor.

$$\text{The LCM} = 2^3 \cdot 3 \cdot 5 \cdot x^3 = 120x^3.$$

$$2. \quad 2x - 4, x^2 - 4, x^2 + 4x + 4$$

$$\left. \begin{aligned} 2x - 4 &= 2 \cdot (x - 2) \\ x^2 - 4 &= (x - 2) \cdot (x + 2) \\ x^2 + 4x + 4 &= (x + 2)^2 \end{aligned} \right\} \text{Complete each polynomial.}$$

Choose the greatest power of each different factor.

$$\text{The LCM} = 2(x - 2)(x + 2)^2.$$

► **Quick check** Find the LCM of $3x + 9$, $x^2 - 9$, $x^2 - 6x + 9$.

We now apply the fundamental principle of rational expressions to “build” a given expression to an equivalent rational expression. To do this, we must **multiply** the numerator and the denominator by the same nonzero polynomial.

■ Example 4-3 C

Write each rational expression as an equivalent expression with the indicated denominator. Assume all denominators are nonzero.

$$1. \quad \frac{13}{3a}, \text{ denominator } 12a^2$$

We obtain the factor with which to build by dividing $\frac{12a^2}{3a} = 4a$.

$$\begin{aligned} \frac{13}{3a} &= \frac{13 \cdot 4a}{3a \cdot 4a} && \text{Multiply the numerator and the denominator by } 4a. \\ &= \frac{52a}{12a^2} && \text{Multiply in the numerator and the denominator.} \end{aligned}$$

$$2. \quad \frac{3x - 2}{2x - 1}, \text{ denominator } 4x^2 - 1$$

Since $4x^2 - 1$ factors to $(2x - 1)(2x + 1)$, the building factor is $2x + 1$.

$$\begin{aligned} \frac{3x - 2}{2x - 1} &= \frac{(3x - 2)(2x + 1)}{(2x - 1)(2x + 1)} && \text{Multiply the numerator and the denominator by } (2x + 1). \\ &= \frac{6x^2 - x - 2}{4x^2 - 1} && \text{Multiply in the numerator and the denominator.} \end{aligned}$$

$$3. \quad \frac{3}{1 - x}, \text{ denominator } x - 1$$

Since $x - 1 = (-1)(1 - x)$, multiply the numerator and the denominator by -1 .

$$\frac{3}{1 - x} = \frac{3 \cdot (-1)}{(1 - x)(-1)} = \frac{-3}{x - 1}$$

► **Quick check** Write $\frac{5z - 3}{3z + 4}$ as an equivalent expression with $9z^2 - 16$ as a denominator. Assume the denominator is nonzero.

We now add and subtract rational expressions that have unlike denominators by building the expressions to equivalent rational expressions having a common denominator—the LCM of the denominators (called the least common denominator, LCD).

Example 4-3 D

Find the indicated sum or difference in lowest terms. Assume no denominator equals zero.

$$1. \frac{6}{5a} + \frac{7}{6b}$$

The LCM of $5a$ and $6b$ is $5a \cdot 6b = 30ab$.

$$\begin{aligned} &= \frac{6}{5a} \cdot \frac{6b}{6b} + \frac{7}{6b} \cdot \frac{5a}{5a} && \text{Building factors} \\ &= \frac{36b}{30ab} + \frac{35a}{30ab} && \text{Multiply by building factors } 6b \text{ and } 5a \\ &= \frac{36b + 35a}{30ab} && \text{Equivalent expressions with LCD } 30ab \\ &= \frac{36b + 35a}{30ab} && \text{Add the numerators.} \end{aligned}$$

$$2. \frac{5}{16a} - \frac{7}{24a^2}$$

$$\left. \begin{aligned} 16a &= 2^4 \cdot a \\ 24a^2 &= 2^3 \cdot 3 \cdot a^2 \end{aligned} \right\} \text{The LCM is } 2^4 \cdot 3 \cdot a^2 = 48a^2.$$

$$\begin{aligned} &= \frac{5}{16a} \cdot \frac{3a}{3a} - \frac{7}{24a^2} \cdot \frac{2}{2} && \text{Building factors} \\ &= \frac{15a}{48a^2} - \frac{14}{48a^2} && \text{Multiply by building factors } 3a \text{ and } 2 \\ &= \frac{15a - 14}{48a^2} && \text{Multiply as indicated} \\ &= \frac{15a - 14}{48a^2} && \text{Subtract the numerators.} \end{aligned}$$

$$3. \frac{10}{x-3} + \frac{9}{x^2+3x-18}$$

$$\left. \begin{aligned} x-3 &= (x-3) \\ x^2+3x-18 &= (x-3)(x+6) \end{aligned} \right\} \text{The LCM is } (x-3)(x+6).$$

$$\begin{aligned} &= \frac{10}{x-3} + \frac{9}{(x-3)(x+6)} && \text{Factor denominators} \\ &= \frac{10}{x-3} \cdot \frac{x+6}{x+6} + \frac{9}{(x-3)(x+6)} && \text{Building factor} \\ &= \frac{10(x+6)}{(x-3)(x+6)} + \frac{9}{(x-3)(x+6)} && \text{Multiply by building factor } x+6 \\ &= \frac{10x+60+9}{(x-3)(x+6)} && \text{Add the numerators} \\ &= \frac{10x+69}{(x-3)(x+6)} && \text{Combine like terms} \end{aligned}$$

$$4. \frac{y+3}{y^2+4y-21} - \frac{y+5}{y^2+6y-27}$$

$$\left. \begin{aligned} y^2+4y-21 &= (y-3)(y+7) \\ y^2+6y-27 &= (y+9)(y-3) \end{aligned} \right\} \text{The LCM is } (y+9)(y-3)(y+7).$$

$$\begin{aligned}
&= \frac{y+3}{(y-3)(y+7)} - \frac{y+5}{(y+9)(y-3)} && \text{Factor denominators} \\
&= \frac{y+3}{(y-3)(y+7)} \cdot \frac{y+9}{y+9} - \frac{y+5}{(y+9)(y-3)} \cdot \frac{y+7}{y+7} && \text{Building factors} \\
&= \frac{(y+3)(y+9)}{(y-3)(y+7)(y+9)} - \frac{(y+5)(y+7)}{(y+9)(y-3)(y+7)} && \text{Multiply in numerator and denominator} \\
&= \frac{(y^2+12y+27)}{(y-3)(y+7)(y+9)} - \frac{(y^2+12y+35)}{(y+9)(y-3)(y+7)} && \text{Multiply in the numerators} \\
&= \frac{(y^2+12y+27) - (y^2+12y+35)}{(y-3)(y+9)(y+7)} && \text{Subtract the numerators} \\
&= \frac{y^2+12y+27-y^2-12y-35}{(y-3)(y+9)(y+7)} && \text{Remove parentheses and change signs} \\
&= \frac{-8}{(y-3)(y+9)(y+7)} && \text{Combine like terms}
\end{aligned}$$

5. $\frac{x^2}{x-8} + \frac{7x+8}{8-x}$

Since the denominators are the opposites of each other, we multiply the numerator and the denominator of $\frac{7x+8}{8-x}$ by -1 .

$$\begin{aligned}
\frac{x^2}{x-8} + \frac{7x+8}{8-x} \cdot \frac{-1}{-1} &= \frac{x^2}{x-8} + \frac{-7x-8}{x-8} && \text{Multiply as indicated} \\
&= \frac{x^2-7x-8}{x-8} && \text{Add numerators} \\
&= \frac{(x+1)(x-8)}{x-8} && \text{Factor the numerator} \\
&= x+1 && \text{Reduce by } x-8
\end{aligned}$$

► **Quick check** Find the difference $\frac{2y+5}{y-2} - \frac{4y+12}{y^2+y-6}$ in lowest terms.

Assume no denominator equals zero.

We now generalize the procedure for adding or subtracting rational expressions.

To add or subtract rational expressions

1. If the denominators are like, add or subtract the numerators and place the result over the common denominator.
2. If the denominators are different, find the least common multiple (LCM) of the denominators.
3. Multiply the numerator and the denominator of each rational expression by all factors present in the LCM but missing in the denominator of the particular rational expression to obtain equivalent rational expressions.
4. Proceed as in step 1. Reduce the results to lowest terms to simplify the answer.

Mastery points**Can you**

- Find the least common multiple (LCM) of a set of polynomials?
- Build rational expressions to equivalent rational expressions?
- Add or subtract rational expressions that have the same denominators?
- Add or subtract rational expressions that have different denominators?

Exercise 4-3

Find the least common multiple of the set of polynomials. See example 4-3 B.

Example $3x + 9, x^2 - 9, x^2 - 6x + 9$

Solution
$$\left. \begin{aligned} 3x + 9 &= 3(x + 3) \\ x^2 - 9 &= (x + 3)(x - 3) \\ x^2 - 6x + 9 &= (x - 3)^2 \end{aligned} \right\} \text{Factor the denominators}$$

The LCM is $3(x + 3)(x - 3)^2$.

1. $14x, 35$

2. $42y, 36$

3. $10k, 16k^2$

4. $18x^3, 21x^2$

5. $32a^3b, 9ab^2$

6. $48x^2y^2, 30x^4y$

7. $xy^3, 6x^2y^2, 15xy^4$

8. $4mn^4, 14m^2n^3, 35mn$

9. $4a, 2a - 4$

10. $9b, 15b + 30$

11. $x - 7, 3x - 21, 6x$

12. $x + 9, 4x + 36, 8x$

13. $p^2 - p - 12, p^2 + 6p + 9, 3p - 12$

14. $n^2 - 9, n^2 - n - 6, 8n^2 + 24n$

15. $a^2 - 25, 10a + 50, 5a - 25$

16. $x^2 + 6x + 9, 2x + 6, x^2 + x - 6$

17. $q^2 - 49, q^2 + 5q - 14, q - 7$

18. $2y + 10, y^2 - 25, y^2 - 10y + 25$

19. $a - b, a^2 - b^2, 5a + 5b$

20. $m - n, 4m + 4n, m^2 - 2mn + n^2$

21. $2a^2 - 13a - 7, 6a^2 + a - 1, a^2 - 49$

22. $4m^2 + 5m - 6, 3m^2 + m - 10, 16m^2 - 9$

Build the given rational expression to equivalent rational expression with the given denominator. Assume all denominators are nonzero. See example 4-3 C.

Example $\frac{5z - 3}{3z + 4}$, denominator $9z^2 - 16$

Solution Since $9z^2 - 16 = (3z + 4)(3z - 4)$, the building factor is $3z - 4$.

$$\begin{aligned} &= \frac{(5z - 3)(3z - 4)}{(3z + 4)(3z - 4)} && \text{Multiply numerator and denominator by } 3z - 4 \\ &= \frac{15z^2 - 29z + 12}{9z^2 - 16} && \text{Multiply in numerator and denominator} \end{aligned}$$

23. $\frac{4}{7x}$, denominator $21x^3$

24. $\frac{8}{9y^2}$, denominator $72y^5$

25. $\frac{5x}{8y}$, denominator $24x^2y^2$

26. $\frac{-3a}{7b^2}$, denominator $35a^2b^3$

27. $4p$, denominator $p - 3$

29. $\frac{p-3}{p+2}$, denominator $p^2 - 4$

31. $\frac{4x}{2x-3}$, denominator $8x^3 - 27$

33. $\frac{2n-1}{n+7}$, denominator $n^2 + 2n - 35$

35. $\frac{2y-5}{4y-1}$, denominator $4y^2 + 7y - 2$

37. $-\frac{3m}{4-5m}$, denominator $25m^2 - 16$

39. $\frac{-9}{9-a}$, denominator $a - 9$

28. $a + 3$, denominator $2a - 1$

30. $\frac{n+7}{n-9}$, denominator $n^2 - 81$

32. $\frac{9x}{3x+1}$, denominator $27x^3 + 1$

34. $\frac{4x+3}{x-9}$, denominator $x^2 - 4x - 45$

36. $\frac{3x+4}{2x-9}$, denominator $8x^2 - 30x - 27$

38. $\frac{9-a}{4-a}$, denominator $a^2 + a - 20$

40. $-\frac{-12}{10-b}$, denominator $b - 10$

Find each indicated sum or difference in lowest terms. Assume all denominators are nonzero. See example 4-3 A.

Example $\frac{4x-3}{x^2-4} - \frac{3x-1}{x^2-4}$

Solution $= \frac{(4x-3) - (3x-1)}{x^2-4}$

Subtract numerators.

$$= \frac{4x-3-3x+1}{x^2-4}$$

Remove parentheses and subtract (change signs).

$$= \frac{x-2}{x^2-4}$$

Combine like terms.

$$= \frac{x-2}{(x-2)(x+2)}$$

Factor the denominator.

$$= \frac{1}{x+2}$$

Reduce by $(x-2)$.

41. $\frac{5}{q} + \frac{8}{q}$

42. $\frac{7}{x} - \frac{17}{x}$

43. $\frac{4y}{y+4} - \frac{7y}{y+4}$

44. $\frac{8a}{a-5} + \frac{7}{a-5}$

45. $\frac{5y}{y+2} + \frac{10}{y+2}$

46. $\frac{3x}{x-4} - \frac{12}{x-4}$

47. $\frac{7x-1}{3x+4} - \frac{4x-5}{3x+4}$

48. $\frac{3y-2}{4y+3} + \frac{5y+8}{4y+3}$

49. $\frac{3x-4}{x^2+5x+6} - \frac{2x-6}{x^2+5x+6}$

50. $\frac{2b+1}{b^2-4} + \frac{1-b}{b^2-4}$

51. $\frac{2x-y}{x+y} - \frac{x-2y}{x+y}$

52. $\frac{3x-2y}{2x+y} - \frac{x+y}{2x+y}$

Find the indicated sum or difference. Assume all denominators are nonzero. See example 4-3 D.

Example $\frac{2y+5}{y-2} - \frac{4y+12}{y^2+y-6}$

Solution $\left. \begin{array}{l} y-2 = (y-2) \\ y^2+y-6 = (y-2)(y+3) \end{array} \right\} \text{Factor denominators}$

The LCM of the denominators is $(y-2)(y+3)$.

$$= \frac{2y+5}{y-2} - \frac{4y+12}{(y-2)(y+3)} \quad \text{Factor denominators}$$

$$= \frac{2y+5}{y-2} \cdot \frac{y+3}{y+3} - \frac{4y+12}{(y-2)(y+3)} \quad \text{Building factor}$$

$$= \frac{(2y+5)(y+3)}{(y-2)(y+3)} - \frac{4y+12}{(y-2)(y+3)} \quad \text{Multiply by building factor}$$

$$= \frac{2y^2+11y+15}{(y-2)(y+3)} - \frac{4y+12}{(y-2)(y+3)} \quad \text{Multiply as indicated}$$

$$= \frac{2y^2+11y+15}{(y-2)(y+3)} - \frac{4y+12}{(y-2)(y+3)}$$

$$= \frac{(2y^2+11y+15) - (4y+12)}{(y-2)(y+3)} \quad \text{Subtract numerators}$$

$$= \frac{2y^2+11y+15-4y-12}{(y-2)(y+3)} \quad \text{Remove parentheses and subtract (change signs)}$$

$$= \frac{2y^2+7y+3}{(y-2)(y+3)} \quad \text{Combine like terms}$$

$$= \frac{(2y+1)(y+3)}{(y-2)(y+3)} \quad \text{Factor numerator}$$

$$= \frac{2y+1}{y-2} \quad \text{Reduce by } (y+3)$$

53. $\frac{8}{3x} + \frac{5}{4x}$

57. $\frac{6}{3x} + \frac{5}{9x^2}$

61. $\frac{4a}{3a+5} + \frac{2a}{2a-3}$

65. $\frac{x+2}{x-9} - \frac{x-6}{9-x}$

69. $\frac{5x}{x^2-2xy-3y^2} - \frac{2y}{x^2+2xy+y^2}$

71. $\frac{a-7}{2a^2+9a-5} + \frac{4-a}{4a^2+23a+15}$

73. $(4a-3) - \frac{2a+5}{5a-2}$

75. $\frac{5}{8p} + \frac{6p-5}{4p^2-8p-60}$

54. $\frac{7}{5a} - \frac{9}{6a}$

58. $\frac{8}{4x^2} - \frac{3}{2x}$

62. $\frac{5m}{m-8} - \frac{3m-8}{2m+7}$

66. $\frac{2a-1}{2a-3} + \frac{4-a}{3-2a}$

55. $\frac{5}{z} - \frac{2}{z^2}$

59. $\frac{2}{3x^2y} - \frac{4}{9xy^2}$

63. $\frac{17}{5y-10} + \frac{19}{2y+4}$

67. $\frac{7}{x^2-5x-6} + \frac{9}{x^2-1}$

70. $\frac{8q}{4q^2-9p^2} + \frac{5q}{4q^2-12pq+9p^2}$

72. $\frac{b+9}{12b^2-5b-2} - \frac{8-2b}{3b^2-17b+10}$

74. $\frac{5x+4}{3x+1} + (8x-5)$

76. $\frac{9p-2}{2p^2-2p-84} + \frac{7}{6p}$

56. $\frac{7}{y^2} + \frac{2}{3y}$

60. $\frac{7}{2ab^2} + \frac{6}{6a^2b}$

64. $\frac{10}{4a-6} - \frac{13}{3a+9}$

68. $\frac{14}{a^2-7a-18} - \frac{8}{a^2-4}$

$$77. \frac{3y}{y^2 + 5y + 6} - \frac{5}{4 - y^2}$$

$$79. \frac{b+3}{b+2} + \frac{2b}{5b^2 - 20}$$

$$81. \frac{2x-3y}{x^2-4xy-12y^2} - \frac{2y-x}{x^2-12xy+36y^2}$$

$$83. \frac{x-3}{8x^2-26x+15} + \frac{3x+2}{6x^2-13x-5}$$

$$85. \frac{5m-n}{8m^2+15mn-2n^2} + \frac{3m+n}{5m^2+6mn-8n^2}$$

$$78. \frac{6a}{6+a-a^2} - \frac{7a}{a^2+7a+10}$$

$$80. \frac{3}{6b^2-4bc} - \frac{4}{9bc-6c^2}$$

$$82. \frac{a-b}{a^2-3ab-4b^2} + \frac{2b-5a}{a^2-16b^2}$$

$$84. \frac{2p-3}{8p^2-18p-5} - \frac{5-7p}{4p^2-27p-7}$$

$$86. \frac{b-2a}{3a^2-2ab-8b^2} - \frac{3a-5b}{9a^2+6ab-8b^2}$$

Solve the following word problems.

87. Workers A , B , C , and D can complete a given job in p , q , r , and s hours, respectively. Working together they can complete in one hour

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} + \frac{1}{s}$$

of the job. Obtain a single expression for what they can do together in one hour.

88. In electricity, the total resistance of any parallel circuit is given by

$$\frac{1}{R_t} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \frac{1}{R_4}$$

Combine the fractions in the right member.

89. Given the focal lengths f_1 and f_2 of two thin lenses that are a distance d apart, the focal length F of the system of lenses is given by

$$\frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2} - \frac{d}{f_1 f_2}$$

Combine the fractions in the right member.

90. In electricity, the true current reading I_t of a current meter is given by

$$I_t = I_r + \frac{R_m}{R_t} \cdot I_r$$

where I_r = the meter reading, R_m = the resistance of the current meter, and R_t = the total resistance of the circuit without the meter. Perform the indicated operations in the right member and write as a single expression in I_r , R_m , and R_t .

91. A lens maker's equation for making a lens is given by

$$\frac{1}{f} = (n-1) \left(\frac{1}{R_1} + \frac{1}{R_2} \right)$$

where f is the focal length of the lens, n is the index of refraction, and R_1 and R_2 are the radii of curvature of the surfaces. Simplify the right member by performing the indicated operations and obtaining a single rational expression.

92. The expression

$$V_1 \left(1 + \frac{T_2 - T_1}{T_1} \right) - V_2 \left(\frac{T_2 - T_1}{T_2} - 1 \right)$$

gives the volume change of a gas under constant pressure. Simplify the expression by performing the indicated operations.

Review exercises

Completely factor the following expressions. See section 3-8.

1. $8x^3 - 2x^2 + 12x - 3$

2. $4x^2 - 12x - 7$

3. $9y^2 - 49$

4. Simplify the expression $4[3 - 2(5 - 1) + 8]$.
See section 1-4.

5. It costs Hank \$24 to rent a mower for one hour. Write a statement for the cost to rent the mower for k hours. See section 2-1.

Student Loans for up to **\$40,000** per year*

Defer payments until after graduation.**
Fast preliminary approval, usually in minutes.

 **Apply Now!**

Go to gmacbankfunding.com

Apply online in as little as 15 minutes

- Loans up to \$40,000 per academic year*
- Good for tuition and other educational expenses: books, fees, a laptop, room and board, travel home, etc.
- Get a check in as few as 5 business days
- Start payments now or up to six months after graduation**
- Reduce your interest rate by as much as 0.50% with automatic payments***

All loans are subject to application and credit approval.

* Undergraduate and graduate borrowers may borrow annually up to the lesser of the cost of attendance or \$30,000 (\$40,000 for certain schools where it has been determined that the annual cost of attendance exceeds \$30,000). Borrowers in the Continuing Education loan program may borrow annually up to \$30,000.

** Undergraduate students may choose to defer repayment until six months after graduation or ceasing to be enrolled at least half time in school. Interest only and immediate repayment options also available.

*** A 0.25% interest rate reduction is available for borrowers who elect to have monthly principal and interest payments transferred electronically from a savings or checking account. The interest rate reduction will begin when automatic principal and interest payments start, and will remain in effect as long as automatic payments continue without interruption. This reduced interest rate will return to contract rate if automatic payments are cancelled, rejected or returned for any reason. Upon request, borrowers are also entitled to an additional 0.25% interest rate reduction if (1) the first 36 payments of principal and interest are paid on time, and (2) at any time prior to the 36th on time payment, the borrower who receives the monthly bill elects to have monthly principal and interest payments transferred electronically from a savings or checking account, and continues to make such automatic payments through the 36th payment. This reduced interest rate will not be returned to contract rate if, after receiving the benefit, the borrower discontinues automatic electronic payment. The lender and servicer reserve the right to modify or discontinue borrower benefit programs (other than the co-signer release benefit) at any time without notice.

GMAC Bank Member FDIC

4-4 ■ Complex rational expressions

The previous sections of this chapter have dealt with simple rational numbers and simple rational expressions—rational numbers and rational expressions that have a single integer, or single polynomial, in the numerator and the denominator. We now consider a **complex rational expression**—a rational expression whose numerator or denominator, or both, contain rational expressions. To illustrate,

$$\frac{\frac{2}{x}}{\frac{3}{x^2}}, \quad \frac{\frac{2}{x} - 3}{4 + \frac{1}{x}}, \quad \frac{\frac{5}{y} - \frac{3}{y}}{y - 3}, \quad \text{and} \quad \frac{\frac{p+1}{p-3}}{\frac{4-p}{p}}$$

are all complex rational expressions. To simplify such expressions, we eliminate the rational expression within the numerator and the denominator to obtain a simple rational expression.

Consider the complex rational expression

$$\left. \begin{array}{l} \frac{p+1}{p-3} \\ \frac{4-p}{p} \end{array} \right\} \begin{array}{l} \text{Primary numerator} \\ \text{Primary denominator} \end{array}$$

For the sake of discussion, we call

$$\frac{p+1}{p-3} \longleftarrow \text{Secondary denominator}$$

the **primary numerator** and

$$\frac{4-p}{p} \longleftarrow \text{Secondary denominator}$$

the **primary denominator**. The expressions $p-3$ and p are called the **secondary denominators**. Thus, to simplify the complex rational expression, we must eliminate the secondary denominators. This can be accomplished in either one of the two ways.

Simplifying a complex rational expression

Method 1 Form a single rational expression in the primary numerator and the primary denominator and divide the primary numerator by the primary denominator.

Method 2 Multiply the primary numerator and the primary denominator by the LCM of the secondary denominators and reduce the resulting fraction to lowest terms.

Example 4-4 A

Simplify each complex rational expression using method 1. Assume all denominators are nonzero.

$$\begin{aligned} 1. \quad \frac{\frac{2}{x}}{\frac{3}{x^2}} &= \frac{2}{x} \div \frac{3}{x^2} \\ &= \frac{2}{x} \cdot \frac{x^2}{3} \\ &= \frac{2x^2}{3x} \\ &= \frac{2x}{3} \end{aligned}$$

Multiply by the reciprocal of $\frac{3}{x^2}$

Multiply numerators and denominators

Reduce by x

$$\begin{aligned} 2. \quad \frac{\frac{2}{x} - 3}{4 + \frac{1}{x}} &= \frac{\frac{2}{x} - \frac{3x}{x}}{\frac{4x}{x} + \frac{1}{x}} = \frac{\frac{2-3x}{x}}{\frac{4x+1}{x}} \\ &= \frac{2-3x}{x} \cdot \frac{x}{4x+1} \\ &= \frac{x(2-3x)}{x(4x+1)} \\ &= \frac{2-3x}{4x+1} \end{aligned}$$

Perform the indicated addition (in the denominator) and the indicated subtraction (in the numerator)

Multiply by the reciprocal of $\frac{4x+1}{x}$

Multiply numerators and denominators

Reduce by x

► **Quick check** Simplify $\frac{\frac{1}{x^2} - \frac{1}{y^2}}{\frac{1}{x} + \frac{1}{y}}$ using method 1. Assume all denominators are nonzero.

A second method to accomplish this same end is to find the least common multiple (LCM) of the *secondary denominators* and to apply the fundamental principle of rational expressions by multiplying the primary numerator and the primary denominator by the LCM, thus eliminating the secondary denominators.

Example 4-4 B

Simplify each complex rational expression using method 2. Assume all denominators are nonzero.

$$1. \frac{\frac{7}{8}}{\frac{5}{6}}$$

The LCM of the secondary denominators, 8 and 6, is 24.

$$\begin{aligned}\frac{7}{8} &= \frac{7 \cdot 3}{8 \cdot 3} \\ \frac{5}{6} &= \frac{5 \cdot 4}{6 \cdot 4} \\ &= \frac{7 \cdot 3}{5 \cdot 4} \\ &= \frac{21}{20}\end{aligned}$$

Multiply the primary numerator and the primary denominator by 24.

Reduce in the numerator and the denominator

Multiply in the numerator and the denominator

$$2. \frac{\frac{p+1}{p-3}}{\frac{4-p}{p}}$$

The LCM of the secondary denominators of $p - 3$ and p is $p(p - 3)$.

$$\begin{aligned}\frac{p+1}{p-3} &= \frac{p+1}{p-3} \cdot \frac{p(p-3)}{p(p-3)} \\ \frac{4-p}{p} &= \frac{4-p}{p} \cdot \frac{p(p-3)}{p(p-3)} \\ &= \frac{(p+1)p}{(4-p)(p-3)} \\ &= \frac{p^2 + p}{-p^2 + 7p - 12}\end{aligned}$$

Multiply the primary numerator and the primary denominator by the LCM $p(p - 3)$.

Reduce in numerator by $p - 3$ and in denominator by p

Multiply in numerator and denominator

$$3. \frac{\frac{5}{y} - \frac{3}{y}}{y - 3}$$

The LCM of the secondary denominators is y .

$$\begin{aligned}\frac{5}{y} - \frac{3}{y} &= \left(\frac{5}{y} - \frac{3}{y} \right) \cdot y \\ &= \frac{5}{y} \cdot y - \frac{3}{y} \cdot y \\ &= \frac{5 - 3}{y(y - 3)} \\ &= \frac{2}{y(y - 3)}\end{aligned}$$

Multiply primary numerator and primary denominator by y .

Apply distributive property

Reduce by y in the numerator

Subtract in the numerator

$$4. \frac{(y+2) + \frac{7}{y-6}}{(y-1) + \frac{4}{y-6}}$$

The LCM of the secondary denominators is $y-6$.

$$\begin{aligned} \frac{(y+2) + \frac{7}{y-6}}{(y-1) + \frac{4}{y-6}} &= \frac{\left[(y+2) + \frac{7}{y-6}\right](y-6)}{\left[(y-1) + \frac{4}{y-6}\right](y-6)} \\ &= \frac{(y+2)(y-6) + \frac{7}{y-6} \cdot (y-6)}{(y-1)(y-6) + \frac{4}{y-6} \cdot (y-6)} \\ &= \frac{y^2 - 4y - 12 + 7}{y^2 - 7y + 6 + 4} \\ &= \frac{y^2 - 4y - 5}{y^2 - 7y + 10} \\ &= \frac{(y-5)(y+1)}{(y-5)(y-2)} \\ &= \frac{y+1}{y-2} \end{aligned}$$

Multiply primary numerator and primary denominator by $y-6$.

Apply distributive property.

Perform indicated operations.

Combine like terms.

Factor numerator and denominator.

Reduce by $y-5$.

$$5. \frac{y^{-1} - x^{-1}}{y^{-1} + x^{-1}}$$

By definition, $y^{-1} = \frac{1}{y}$ and $x^{-1} = \frac{1}{x}$. Thus,

$$\frac{y^{-1} - x^{-1}}{y^{-1} + x^{-1}} = \frac{\frac{1}{y} - \frac{1}{x}}{\frac{1}{y} + \frac{1}{x}}$$

The LCM of the secondary denominators is xy .

$$\begin{aligned} &= \frac{\frac{1}{y} \cdot xy - \frac{1}{x} \cdot xy}{\frac{1}{y} \cdot xy + \frac{1}{x} \cdot xy} \\ &= \frac{x - y}{x + y} \end{aligned}$$

Multiply the primary numerator and primary denominator by xy using the distributive property.

Reduce in each term.

► **Quick check** Simplify $\frac{\frac{x-3}{x+2}}{\frac{x+1}{x-4}}$ using method 2. Assume all denominators are nonzero.

Note A reminder again—always check the factorability of the numerator and the denominator to *reduce* the simple rational expression to lowest terms.

Mastery points*Can you*

- Recognize a complex rational expression?
- Simplify a complex rational expression?

Exercise 4-4

Simplify each complex rational expression using method 1. Assume all denominators are nonzero. See example 4-4 A.

Example
$$\frac{\frac{1}{x^2} - \frac{1}{y^2}}{\frac{1}{x} + \frac{1}{y}}$$

Solution
$$= \frac{\frac{y^2 - x^2}{x^2 y^2}}{\frac{y + x}{xy}}$$

$$= \frac{\frac{xy}{x^2 y^2} \cdot \frac{xy}{y + x}}{\frac{(y - x)(y + x)xy}{x^2 y^2 (y + x)}}$$

$$= \frac{y - x}{xy}$$

Subtract in the numerator and add in the denominator

Multiply by the reciprocal of $\frac{y + x}{xy}$

Factor and multiply numerators and denominators

Reduce by common factors xy and $y + x$

1. $\frac{\frac{3}{5}}{\frac{4}{7}}$

2. $\frac{\frac{5}{6}}{\frac{7}{8}}$

3. $\frac{\frac{2}{3} + \frac{1}{4}}{\frac{1}{2} - \frac{1}{4}}$

4. $\frac{\frac{1}{3} + \frac{1}{2}}{\frac{2}{3} - \frac{5}{6}}$

5. $\frac{4 - \frac{2}{5}}{3 + \frac{3}{10}}$

6. $\frac{\frac{4}{x}}{\frac{8}{x^2}}$

7. $\frac{\frac{7}{m^2}}{\frac{8}{m}}$

8. $\frac{\frac{5}{b}}{\frac{9}{b - 3}}$

9. $\frac{\frac{-4}{y + 1}}{\frac{3}{y}}$

10. $\frac{\frac{8}{a - 3}}{\frac{-6}{a + 2}}$

11. $\frac{\frac{x}{x + 7}}{\frac{x}{x - 3}}$

12. $\frac{\frac{m - 3}{4}}{\frac{2m + 7}{6}}$

13. $\frac{\frac{1}{a} + \frac{1}{b}}{\frac{1}{a^2} - \frac{1}{b^2}}$

14. $\frac{\frac{x - y}{1}}{\frac{1}{x} - \frac{1}{y}}$

15. $\frac{\frac{x}{y} + 2}{\frac{x}{y} - \frac{4y}{x}}$

16. $\frac{\frac{2}{q} - \frac{3}{r}}{2 + \frac{1}{qr}}$

Simplify each complex rational expression by multiplying the numerator and the denominator by the LCM of the secondary denominators. See example 4-4 B.

Example
$$\frac{\frac{x-3}{x+2}}{\frac{x+1}{x-4}}$$

Solution
$$\begin{aligned} &= \frac{\frac{x-3}{x+2} \cdot (x+2)(x-4)}{\frac{x+1}{x-4} \cdot (x+2)(x-4)} \\ &= \frac{(x-3)(x-4)}{(x+1)(x+2)} \\ &= \frac{x^2 - 7x + 12}{x^2 + 3x + 2} \end{aligned}$$

Multiply primary numerator and primary denominator by LCM $(x+2)(x-4)$

Reduce by common factors

Perform indicated multiplication

17.
$$\frac{\frac{5}{9}}{\frac{3}{4}}$$

18.
$$\frac{\frac{1}{6} - \frac{7}{8}}{\frac{2}{3} + \frac{1}{4}}$$

19.
$$\frac{\frac{5}{y^2}}{\frac{7}{y}}$$

20.
$$\frac{\frac{3}{x-2}}{\frac{9}{x}}$$

21.
$$\frac{\frac{7}{a+9}}{\frac{-4}{a-5}}$$

22.
$$\frac{\frac{n-9}{n+2}}{\frac{n}{3n-5}}$$

23.
$$\frac{\frac{4x^2 - y^2}{3x}}{2x + y}$$

24.
$$\frac{\frac{x+3}{y}}{\frac{x^2 - 9y^2}{6y}}$$

25.
$$\frac{4 - \frac{3}{x+3}}{5 + \frac{6}{x-1}}$$

26.
$$\frac{\frac{7}{y-3} + 8}{9 - \frac{1}{2y+3}}$$

27.
$$\frac{\frac{1}{x} + \frac{1}{y^2}}{\frac{1}{x^2} - \frac{1}{x}}$$

28.
$$\frac{\frac{1}{a} - \frac{1}{b}}{\frac{1}{a^2} + \frac{1}{b^2}}$$

29.
$$\frac{\frac{m+n}{1}}{\frac{1}{m} - \frac{1}{n}}$$

30.
$$\frac{\frac{x-y}{1}}{\frac{1}{x} + \frac{1}{y}}$$

31.
$$\frac{\frac{5}{p^2} + \frac{4}{q}}{p - q}$$

32.
$$\frac{\frac{1}{m^2} - \frac{1}{n^2}}{m + n}$$

33.
$$\frac{(a+5) + \frac{3}{a+4}}{(a+3) - \frac{5}{a+4}}$$

34.
$$\frac{(x-3) + \frac{7}{2x+1}}{(x+9) - \frac{3}{2x+1}}$$

35.
$$\frac{y - \frac{3}{4y-3}}{(y+2) + \frac{3}{y+5}}$$

36.
$$\frac{(m-3) + \frac{6}{2m+3}}{m - \frac{9}{m-6}}$$

37.
$$\frac{\frac{t^2 - 2t - 8}{t^2 + 7t + 6}}{\frac{t^2 - t - 6}{t^2 + 2t + 1}}$$

38.
$$\frac{\frac{y^2 - 5y - 14}{y^2 + 3y - 10}}{\frac{y^2 - 8y + 7}{y^2 + 6y + 5}}$$

39.
$$\frac{\frac{3}{x^2 - x - 6}}{\frac{2}{x+2} - \frac{4}{x-3}}$$

40.
$$\frac{\frac{9}{a-7} + \frac{8}{2a+3}}{\frac{10}{2a^2 - 11a - 21}}$$

41.
$$\frac{\frac{5}{7} + \frac{4}{b-1}}{\frac{7}{b+5} - \frac{3}{b-1}}$$

42.
$$\frac{\frac{6}{x+5} - 7}{\frac{8}{x+5} - \frac{9}{x+3}}$$

43.
$$\frac{\frac{1}{1-x} - \frac{1}{1+x}}{\frac{1}{1+x} - \frac{1}{1-x}}$$

$$44. \frac{\frac{a-1}{a-b} - \frac{a-1}{a-1}}{\frac{1}{a-1} - \frac{1}{a-b}}$$

$$48. \frac{x^2 - y^2}{x^{-1} + y^{-1}}$$

$$52. \frac{2x^{-1} - y^{-1}}{x^{-1} + 5y^{-1}}$$

$$45. \frac{\frac{3}{ab} + \frac{4}{bc} - \frac{2}{ac}}{\frac{5}{abc}}$$

$$49. \frac{x^{-1}}{x^{-1} + y^{-1}}$$

$$53. (p^{-1} - q^{-1})^{-1}$$

$$46. \frac{\frac{x}{yz} - \frac{y}{xz} + \frac{z}{xy}}{\frac{1}{x^2y^2} - \frac{1}{x^2z^2} + \frac{1}{y^2z^2}}$$

$$50. \frac{x^{-2}}{x^{-2} - y^{-2}}$$

$$54. (x^{-1} + y^{-1})^{-1}$$

$$47. \frac{x^{-1} + y^{-1}}{x^{-1} - y^{-1}}$$

$$51. \frac{p^{-2} + q^{-2}}{q^{-2}}$$

Solve the following word problems.

55. In electricity, a relationship for the current, i , in a capacitor of “size” C is given by

$$i = \frac{V_s - \frac{it}{C}}{R}$$

Simplify the right member.

56. In electricity, the voltage between two adjacent nodes, denoted by $V_{AA'}$, is given by Millman's Theorem,

$$V_{AA'} = \frac{\frac{V_{S1}}{R_1} + \frac{V_{S2}}{R_2} + \frac{V_{S3}}{R_3}}{\frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}}$$

where V_{S1} , V_{S2} , V_{S3} represent equivalent voltages of the branches between nodes A and A' , whereas R_1 , R_2 , and R_3 are equivalent resistances of the branches between nodes A and A' . Simplify the right member.

57. A refrigeration coefficient, CP , of performance formula for the ideal refrigerator is given by

$$CP = \frac{1}{\frac{T_2}{T_1} - 1}$$

Simplify the right member.

58. The capacitance C of a circuit connecting three capacitances C_1 , C_2 , and C_3 in series is given by

$$C = \frac{1}{\frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3}}$$

Simplify the right member.

59. When making a round trip whose one-way distance is d , the average rate (speed) traveled, r , is given by

$$r = \frac{2d}{\frac{d}{r_1} + \frac{d}{r_2}}$$

where r_1 is the average rate going and r_2 is the average rate coming back. Simplify the right member.

Review exercises

Find the solution set of the following equations. See sections 2-1 and 2-4.

$$1. \frac{1}{2}x + 3 = \frac{2}{3}$$

$$2. |x - 3| = 5$$

$$3. 5(2x - 4) + 3x = 2x + 2$$

Find the solution set of the following inequalities. See section 2-6.

$$4. |x - 2| < 4$$

$$5. |2x + 5| \geq 3$$

6. Perform the indicated subtraction. See section 3-2.

$$(3a^3 + 2a^2 - 3) - (a^3 - 6a^2 + a + 1)$$

Campfire queen Cycling champion Sentimental geologist*

Learn more about
Marjon Walrod
and tell us more
about you. Visit
pwc.com/bringit.

Your life. You can
bring it with you.



*connectedthinking

PRICEWATERHOUSECOOPERS 

© 2006 PricewaterhouseCoopers LLP. All rights reserved. "PricewaterhouseCoopers" refers to PricewaterhouseCoopers LLP (a Delaware limited liability partnership) or, as the context requires, the PricewaterhouseCoopers global network or other member firms of the network, each of which is a separate and independent legal entity. *connectedthinking is a trademark of PricewaterhouseCoopers LLP (US). We are proud to be an Affirmative Action and Equal Opportunity Employer.

4-5 ■ Quotients of polynomials

Division of a polynomial by a monomial

The first type of division that we will study is that of a polynomial divided by a monomial. Consider the indicated division

$$\frac{4a^3 - 12a^2 + 8a}{2a}$$

Using the meaning of division, we can rewrite this as

$$(4a^3 - 12a^2 + 8a) \cdot \frac{1}{2a}$$

and applying the distributive property, we have

$$\begin{aligned}\frac{4a^3 - 12a^2 + 8a}{2a} &= \frac{4a^3}{2a} - \frac{12a^2}{2a} + \frac{8a}{2a} \\ &= 2a^2 - 6a + 4\end{aligned}$$

Dividing a polynomial by a monomial

For monomials a_1, a_2, \dots, a_n and d , where $d \neq 0$,

$$\frac{a_1 + a_2 + \dots + a_n}{d} = \frac{a_1}{d} + \frac{a_2}{d} + \dots + \frac{a_n}{d}$$

Concept

To divide a polynomial by a monomial, simply divide each term of the polynomial by the monomial and write the resulting quotients.

■ **Example 4-5 A**

Perform the indicated division. Assume that no denominator equals zero.

- $$\frac{4a^6 + 16a^4 - 6a}{4a^2} = \frac{4a^6}{4a^2} + \frac{16a^4}{4a^2} - \frac{6a}{4a^2}$$

Divide each term by monomial $4a^2$

$$= a^4 + 4a^2 - \frac{3}{2a}$$

Reduce to lowest terms
- $$\frac{x(3y - 2) + 2(3y - 2)}{3y - 2} = \frac{x(3y - 2)}{(3y - 2)} + \frac{2(3y - 2)}{(3y - 2)}$$

Divide each term by $(3y - 2)$

$$= x + 2$$

Reduce to lowest terms

Note A common error in this type of problem is demonstrated in the example

$$\frac{x^3 + x^2}{x^2} = \frac{x^3 + \cancel{x^2}}{\cancel{x^2}} = \frac{x^3 + 1}{1} = x^3 + 1$$

It is tempting to "cancel" the x^2 in the numerator with the x^2 in the denominator, but the correct procedure is

$$\frac{x^3 + x^2}{x^2} = \frac{x^3}{x^2} + \frac{x^2}{x^2} = x + 1$$

► **Quick check** Simplify $\frac{5x^3 - 10x^2 + 25x}{5x^2}$. Assume the denominator is nonzero.

Division of a polynomial by a polynomial

The second type of division that we will study is that of a polynomial divided by another polynomial. Consider the example

$$\frac{2x^2 - x - 15}{x - 3}$$

The example is set up so that the divisor, $x - 3$, and the dividend, $2x^2 - x - 15$, are arranged in descending powers of the variable.

$$x - 3 \overline{) 2x^2 - x - 15}$$

Now we divide the first term of the dividend, $2x^2$, by the first term of the divisor, x . The result is $\frac{2x^2}{x} = 2x$, and we place the $2x$ above the division line.

$$x - 3 \overline{) 2x^2 - x - 15} \quad \begin{array}{r} 2x \\ \hline \end{array}$$

Next, we multiply $x - 3$ and $2x$, placing the product below the dividend, and subtract.

$$x - 3 \overline{) 2x^2 - x - 15} \quad \begin{array}{r} 2x \\ \hline 2x^2 - 6x \\ \hline \end{array} \quad 2x(x - 3) = 2x^2 - 6x$$

Recall that when we subtract, we change the signs and add. Then $(2x^2 - x) - (2x^2 - 6x) = 2x^2 - x - 2x^2 + 6x = 5x$.

$$x - 3 \overline{) 2x^2 - x - 15} \quad \begin{array}{r} 2x \\ \hline -2x^2 - 6x \\ \hline 5x \\ \hline \end{array}$$

Now we bring down the -15 and repeat the same process.

$$x - 3 \overline{) 2x^2 - x - 15} \quad \begin{array}{r} 2x \\ \hline 2x^2 - 6x \\ \hline 5x - 15 \\ \hline \end{array}$$

Divide the $5x$ by x , which results in 5 , and multiply 5 and $x - 3$.

$$x - 3 \overline{) 2x^2 - x - 15} \quad \begin{array}{r} 2x + 5 \\ \hline 2x^2 - 6x \\ \hline 5x - 15 \\ \hline 5x - 15 \\ \hline \end{array}$$

Subtract $(5x - 15) - (5x - 15) = 5x - 15 - 5x + 15 = 0$.

$$x - 3 \overline{) 2x^2 - x - 15} \quad \begin{array}{r} 2x + 5 \\ \hline 2x^2 - 6x \\ \hline 5x - 15 \\ \hline 5x - 15 \\ \hline 0 \end{array}$$

We subtract and get a remainder of zero, so the quotient is $2x + 5$.

$$\frac{2x^2 - x - 15}{x - 3} = 2x + 5$$

To check the problem, multiply the quotient times the divisor to get the dividend.

$$(2x + 5)(x - 3) = 2x^2 - x - 15$$

Note It is important when dividing a polynomial by a polynomial that both the divisor and the dividend have their terms arranged in descending powers of the variable.

Example 4-5 B

Find the indicated quotients. Assume no divisor is equal to zero.

1. $(2x^2 - 11x + 15) \div (x - 3)$

$$\begin{array}{r} 2x - 5 \\ x - 3 \overline{) 2x^2 - 11x + 15} \\ \underline{2x^2 - 6x} \\ -5x + 15 \\ \underline{-5x + 15} \\ 0 \end{array}$$

Subtract (change signs and add)

Subtract (change signs and add)

Therefore $\frac{2x^2 - 11x + 15}{x - 3} = 2x - 5$.

2. $(2y^3 - y^2 + 5y + 5) \div (2y + 1)$

$$\begin{array}{r} y^2 - y + 3 \\ 2y + 1 \overline{) 2y^3 - y^2 + 5y + 5} \\ \underline{2y^3 + y^2} \\ -2y^2 + 5y \\ \underline{-2y^2 - y} \\ 6y + 5 \\ \underline{6y + 3} \\ 2 \end{array}$$

Subtract (change signs and add)

Subtract (change signs and add)

Subtract (change signs and add)

Remainder

When there is a remainder, as in this example, we write the remainder over the divisor.

$$\frac{2y^3 - y^2 + 5y + 5}{2y + 1} = y^2 - y + 3 + \frac{2}{2y + 1}$$

3. $(2x^3 - x + 5) \div (x - 1)$

There is no term in the dividend that contains x^2 . Therefore we will insert $0x^2$ as a placeholder so that all powers of the variable x are present in descending order. The value of the dividend has not changed since we added $0x^2$, which is another name for 0.

$$\begin{array}{r} 2x^2 + 2x + 1 \\ x - 1 \overline{) 2x^3 + 0x^2 - x + 5} \\ \underline{2x^3 - 2x^2} \\ 2x^2 - x \\ \underline{2x^2 - 2x} \\ x + 5 \\ \underline{x - 1} \\ 6 \end{array}$$

Subtract (change signs and add)

Subtract (change signs and add)

Subtract (change signs and add)

Remainder

Hence $\frac{2x^3 - x + 5}{x - 1} = 2x^2 + 2x + 1 + \frac{6}{x - 1}$.

► **Quick check** Divide $\frac{3y^3 + 5y^2 - 11y + 6}{3y - 1}$. Assume the denominator is nonzero.

Mastery points

Can you

- Divide a polynomial by a monomial?
- Divide a polynomial by a polynomial?

Exercise 4-5

Perform the indicated divisions. Assume that no divisor is equal to zero. See example 4-5 A.

Example $\frac{5x^3 - 10x^2 + 25x}{5x^2}$

Solution
$$\begin{aligned} &= \frac{5x^3}{5x^2} - \frac{10x^2}{5x^2} + \frac{25x}{5x^2} \\ &= x - 2 + \frac{5}{x} \end{aligned}$$

Divide denominator into each term

Divide in each term by common factors

1. $\frac{25x^2 - 15x + 10}{5}$

2. $\frac{2a^4 - 3a^2 + a}{a}$

3. $\frac{4x^4 - 8x^3 + 12x}{-4x}$

4. $\frac{15y^5 + 25y^2 + 10y}{-5y^2}$

5. $\frac{ac^2 - ac}{ac}$

6. $\frac{bx - b^2x^2}{bx}$

7. $\frac{30x^3y^4 + 21x^2y^2 - 18x^2y^4}{3x^2y^2}$

8. $\frac{36x^4y^2z^3 - 24x^2y^5z + 18x^2y^2z^2}{6x^2y^3z}$

9. $\frac{21a^7b^2c^3 - 35a^5b^5c^3 + 49a^4b^2c^3}{7abc^3}$

10. $\frac{x(y - 2) - z(y - 2)}{y - 2}$

11. $\frac{2a(b - 4) + 3c(b - 4)}{b - 4}$

12. $\frac{x^2y(z + 3) - 3x^4y^3(z + 3)}{xy(z + 3)}$

See example 4-5 B.

Example $\frac{3y^3 + 5y^2 - 11y + 6}{3y - 1}$

Solution

$$\begin{array}{r} y^2 + 2y - 3 \\ 3y - 1 \overline{) 3y^3 + 5y^2 - 11y + 6} \\ \underline{3y^3 - y^2} \downarrow \\ 6y^2 - 11y \downarrow \\ \underline{6y^2 - 2y} \downarrow \\ -9y + 6 \downarrow \\ \underline{-9y + 3} \\ 3 \end{array}$$

Subtract (change signs and add)

Subtract (change signs and add)

Subtract (change signs and add)

Remainder

$$\frac{3y^3 + 5y^2 - 11y + 6}{3y - 1} = y^2 + 2y - 3 + \frac{3}{3y - 1}$$

13. $(a^2 - 2a - 8) \div (a + 2)$

16. $(2x - 5 + x^2) \div (x + 4)$

19. $\frac{a^3 + a^2 - 2a + 12}{a + 3}$

22. $\frac{3x^3 - 4x^2 - 5x - 4}{x + 2}$

25. $\frac{4a^3 + 8a^2 - 5a + 1}{2a + 1}$

28. $\frac{27x^3 - 1}{3x - 1}$

31. $\frac{3x^4 - 2x^3 + x - 1}{x + 1}$

34. $\frac{a^4 - 2a^2 - 3a - 1}{a^2 - 2a - 1}$

37. $\frac{x^4 + 4x^3 + x^2 - 10x - 12}{x^2 + x - 4}$

40. $\frac{a^4 + 4a^3 + a^2 - 10a - 9}{a^2 + 3a + 2}$

14. $(y^2 + 2y - 3) \div (y + 3)$

17. $(10 - 7x + x^2) \div (x - 3)$

20. $\frac{y^3 + 3y^2 - y - 6}{y + 2}$

23. $\frac{5x^2 - 11x + 2x^3 + 4}{2x - 1}$

26. $\frac{6y^3 - y^2 - 11y + 10}{3y - 2}$

29. $\frac{a^3 + 27}{a + 3}$

32. $\frac{2x^3 + 5x^2 + 5x + 3}{x^2 + x + 1}$

35. $\frac{y^4 + 4y^3 + 3y^2 - 2y - 1}{y^2 + 3y + 1}$

38. $\frac{2x^4 - x^3 + 5x^2 - x + 3}{x^2 + 1}$

41. $\frac{2y^4 - 3y^3 + 8y^2 - 9y + 8}{2y^2 - 3y + 2}$

15. $(y^2 + 7y + 11) \div (y + 5)$

18. $\frac{a^2 - 5a + 1}{a - 1}$

21. $\frac{2y^3 - y^2 - 2y + 3}{y + 1}$

24. $\frac{3a^3 - 2 - 5a - a^2}{3a + 2}$

27. $\frac{9y^3 + 11y + 6}{3y + 1}$

30. $\frac{x^4 - 16}{x - 2}$

33. $\frac{3a^3 - 4a^2 + 10a - 3}{a^2 - a + 3}$

36. $\frac{y^4 - y^3 - 11y^2 + 10y + 2}{y^2 - 4y + 2}$

39. $\frac{3a^4 - 2a^3 + 2a^2 - 4a - 8}{a^2 + 2}$

42. $\frac{3y^4 + y^3 - 8y^2 - 3y - 5}{3y^2 + y + 1}$

Solve the following word problems.

43. Evaluate
- $2y^3 - y^2 - 2y + 3$
- when
- $y = -1$
- and compare this answer to the remainder found in exercise 21.

44. Evaluate
- $3x^3 - 4x^2 - 5x - 4$
- when
- $x = -2$
- and compare this answer to the remainder found in exercise 22.

45. The area of a rectangle is found by multiplying the length times the width. If the area of a rectangle is
- $6x^2 - 17x + 12$
- and the length is
- $3x - 4$
- , find the width.

46. A contractor uses the expression
- $x^2 + 6x + 8$
- to represent the square footage of a room. If she decides that the length of the room will be represented by
- $x + 4$
- , what will the width of the room be in terms of
- x
- ?

47. The volume of a box is found by multiplying the length times the width times the height. If the volume of a box is
- $6x^3 + 11x^2 - 19x + 6$
- , the height is
- $x + 3$
- , and the width is
- $2x - 1$
- , find the length.

48. An electrician uses the expression
- $x^2 + 5x + 6$
- to determine the amount of wire to order when wiring a house. If the formula comes from multiplying the number of rooms times the number of outlets and he knows the number of rooms to be
- $x + 2$
- , find the number of outlets in terms of
- x
- .

Looking for more **FREE** Books?

Then visit the web's home for free textbooks:

www.TextbookRevolution.org

At [Textbook Revolution](http://www.TextbookRevolution.org), you'll find links to textbooks and select educational resources of all kinds. ***All** of the books are offered for **free** online viewing by their respective copyright holders.*

Textbook Revolution has links to hundreds of books—math books, science books, business books, and much more. The site is non-profit and student-run. There's no sign-up and no fee. Ask your professors to join the Textbook Revolution and assign free books from www.TextbookRevolution.org!

College is expensive enough! Join the revolution—the [Textbook Revolution](http://www.TextbookRevolution.org)—and take the bite out of your books.

Review exercises

Simplify the following expressions. Assume all denominators are nonzero. Answer with positive exponents only. See sections 3-1 and 3-3.

1. $\frac{x^3y}{x^2y}$

2. $\frac{x^{-3}}{x^{-6}}$

3. $(2a^{-4}b^0)^3$

4. Simplify the expression $\frac{\frac{3}{4} - \frac{1}{2}}{\frac{2}{3} + \frac{1}{6}}$. See section 4-4.

Perform the indicated operations. Assume all denominators are nonzero. See sections 4-2 and 4-3.

5. $\frac{2x-1}{x^2-9} - \frac{x+1}{x+3}$

6. $\frac{2x-1}{3x-1} \div \frac{4x}{6x-2}$

7. Find the solution set of the equation $\frac{1}{5}x - \frac{2}{3} = \frac{3}{5}$.
See section 2-1.

4-6 ■ Synthetic division, the remainder theorem, and the factor theorem

In section 4-5, we studied the procedure for dividing a polynomial by another polynomial. Many times the divisor is a binomial of the form $x - k$, k is a constant. We can shorten the process considerably by using **synthetic division**.

Consider the following example: $(2x^3 + 5x^2 + x - 1) \div (x + 2)$.
Performing the indicated division, we obtain

$$\begin{array}{r}
 \text{Divisor } x + 2 \overline{) 2x^3 + 5x^2 + x - 1} \quad \begin{array}{l} \text{Quotient} \\ \text{Dividend} \end{array} \\
 \underline{2x^3 + 4x^2} \\
 x^2 + x \\
 \underline{x^2 + 2x} \\
 -x - 1 \\
 \underline{-x - 2} \\
 1 \quad \text{Remainder}
 \end{array}$$

We can write the exact same problem using only the coefficients of the terms.

$$\begin{array}{r}
 \begin{array}{l} \text{Coefficients from} \\ \text{the divisor} \rightarrow 1 \end{array} \quad \begin{array}{r} 2 \quad 1 \quad -1 \quad -1 \\ 2 \overline{) 2 \quad 5 \quad 1 \quad -1} \\ \underline{2 \quad 4} \\ 1 \quad 1 \\ \underline{1 \quad 2} \\ -1 \quad -1 \\ \underline{-1 \quad -2} \\ 1 \end{array} \quad \begin{array}{l} \text{Coefficients from the quotient} \\ \text{Coefficients from the dividend} \end{array} \\
 \text{Remainder}
 \end{array}$$

Observe that the circled numbers are repetitions of the numbers directly above them. We can rewrite the problem without them.

$$\begin{array}{r|rrrr}
 1 & 2 & 5 & 1 & -1 \\
 & & 4 & & \\
 & & 1 & \textcircled{1} & \\
 & & & 2 & \\
 & & & -1 & \textcircled{-1} \\
 & & & & -2 \\
 & & & & 1
 \end{array}$$

The circled numbers are again the same as the numbers directly above them. Therefore we can omit them.

$$\begin{array}{r|rrrr}
 1 & 2 & 5 & 1 & -1 \\
 & & 4 & & \\
 & & 1 & & \\
 & & & 2 & \\
 & & & -1 & \\
 & & & & -2 \\
 & & & & 1
 \end{array}$$

All of the numbers can be condensed. We omit the top row of numbers since it duplicates the bottom set of numbers. We shall also omit the 1 at the upper left.

$$\begin{array}{r|rrrr}
 2 & 5 & 1 & -1 \\
 & 4 & 2 & -2 \\
 \hline
 2 & 1 & -1 & 1
 \end{array}$$

By changing the 2 at the upper left to its additive inverse, -2 , and adding the additive inverse in each step, instead of subtracting, we obtain the same result. The following is the final form to be used when performing synthetic division.

$$\begin{array}{r|rrrr}
 -2 & 2 & 5 & 1 & -1 \\
 & & -4 & -2 & 2 \\
 \hline
 2 & 1 & -1 & 1 \\
 \uparrow & \uparrow & \uparrow & \uparrow \\
 x^2 & x & \text{Constant} & \text{Remainder}
 \end{array}$$

The bottom line of this synthetic division process represents the coefficients of the terms of the quotient along with the remainder. That is, $2 \quad 1 \quad -1 \quad 1$ represents the quotient $2x^2 + x - 1$ with a remainder of 1.

Note The degree 2 of the quotient is one less than the degree 3 of the dividend. This will *always* be the case.

Example 4-6 A

Divide the following using synthetic division. Assume no denominator equals zero.

1. $(3x^2 + 7x + 6) \div (x + 1)$

To use synthetic division, the divisor *must* be of the form $x - k$. Therefore we write $x + 1$ as $x - (-1)$ so $k = -1$. Now write the coefficients of the terms in $3x^2 + 7x + 6$ with -1 to the left.

$$\begin{array}{r|rrr} -1 & 3 & 7 & 6 \end{array}$$

First, we bring down 3 and multiply that by -1 . This product, -3 , is added to 7.

$$\begin{array}{r|rrr} -1 & 3 & 7 & 6 \\ \downarrow & & -3 & \\ \hline & 3 & 4 & \end{array}$$

The 4 is now multiplied by -1 and the product, -4 , is added to 6.

$$\begin{array}{r|rrr} -1 & 3 & 7 & 6 \\ & & -3 & -4 \\ \hline & 3 & 4 & 2 \end{array}$$

We can now read the coefficients of the quotient and the remainder from the last row. The answer is $3x + 4 + \frac{2}{x + 1}$. (2 is the remainder.)

2. $(2x^3 - 7x^2 - x + 12) \div (x - 3)$

The divisor, $x - 3$, is already in the form $x - k$, where $k = 3$. We set up the problem as follows:

$$\begin{array}{r|rrrr} 3 & 2 & -7 & -1 & 12 \end{array}$$

We bring down 2 and begin the repetitive process of multiplying the last number in the bottom row times k , that is, 3, and adding this to the value in the next column.

$$\begin{array}{r|rrrr} 3 & 2 & -7 & -1 & 12 \\ \downarrow & & 6 & -3 & -12 \\ \hline & 2 & -1 & -4 & 0 \end{array}$$

Since we have a zero in the last position of the bottom row, there is no remainder and the quotient is $2x^2 - x - 4$.

3. $(x^3 - 11x + 8) \div (x - 3)$

Since the x^2 term is missing in the dividend, we think of the expression as $x^3 + 0x^2 - 11x + 8$ when writing down the coefficients.

$$\begin{array}{r|rrrr} 3 & 1 & 0 & -11 & 8 \\ \downarrow & & 3 & 9 & -6 \\ \hline & 1 & 3 & -2 & 2 \end{array}$$

The answer is then $x^2 + 3x - 2 + \frac{2}{x - 3}$.

$$4. (2x^4 - 10x^3 + 11x^2 - 11x + 8) \div (x - 2)$$

$$\begin{array}{r|rrrrr} 2 & 2 & -10 & 11 & -11 & 8 \\ & \downarrow & & & & \\ & 4 & -12 & -2 & -26 & \\ \hline & 2 & -6 & -1 & -13 & -18 \end{array}$$

The answer is $2x^3 - 6x^2 - x - 13 - \frac{18}{x-2}$.

► **Quick check** Use synthetic division to divide $(x^3 + 10x - 2) \div (x + 2)$. ■

Remainder and factor theorems

From our work with synthetic division, we can see that division by a polynomial in x , $P(x)$, by a polynomial $x - r$ results in a quotient $Q(x)$ and a constant remainder R . That is,

$$\frac{P(x)}{x - r} = Q(x) + R$$

and so

$$P(x) = (x - r)Q(x) + R$$

Using this equation, we evaluate $P(r)$ when $x = r$.

$$\begin{aligned} P(x) &= (x - r)Q(x) + R \\ P(r) &= (r - r)Q(r) + R \\ &= 0 \cdot Q(r) + R \\ &= 0 + R \\ &= R \end{aligned}$$

Thus, $P(r) = R$ and we find that the value of the polynomial when $x = r$ is the remainder R . We have just proved the **Remainder Theorem**.

Remainder theorem

If a polynomial $P(x)$ is divided by $x - r$, where r is a real number, the remainder R is $P(r)$.

■ Example 4-6 B

- Determine the remainder when $P(x) = 2x^3 - 5x^2 + 4x - 2$ is divided by $x - 3$. Evaluate $P(3)$ using substitution.
 - Using synthetic division,

$$\begin{array}{r|rrrr} 3 & 2 & -5 & 4 & -2 \\ & \downarrow & & & \\ & 6 & 3 & 21 & \\ \hline & 2 & 1 & 7 & 19 \end{array} \quad \leftarrow \text{Remainder}$$

Thus, $P(3) = 19$

b. Using substitution, we must show that $P(3) = 19$.

$$\begin{aligned} P(x) &= 2x^3 - 5x^2 + 4x - 2 \\ P(3) &= 2(3)^3 - 5(3)^2 + 4(3) - 2 \\ &= 54 - 45 + 12 - 2 \\ &= 19 \end{aligned}$$

Thus, we have shown that $R = 19$ is the result in either case.

► **Quick check** Using synthetic division and the remainder theorem, find the remainder when $P(x) = 3x^3 - 2x^2 + 4x - 5$ is divided by $x + 4$. Evaluate $P(-4)$ using substitution.

The **factor theorem**, which is a direct result of the remainder theorem (a corollary), applies when the remainder $R = 0$. This theorem shows there is a close relationship between the *factors* of a polynomial $P(x)$ and the values of x for which $P(x) = 0$.

Factor theorem

Given polynomial $P(x)$ and real number r , then

$x - r$ is a factor of $P(x)$ if and only if $P(r) = 0$

Example 4-6 C

1. Show that $x + 2$ is a factor of $P(x) = x^3 - 3x^2 - 6x + 8$.

By the factor theorem, $x + 2$ is a factor of $P(x)$ if $P(-2) = 0$. Using synthetic division, we show that the remainder $R = 0$.

$$\begin{array}{r|rrrr} -2 & 1 & -3 & -6 & 8 \\ & \downarrow & -2 & 10 & -8 \\ \hline & 1 & -5 & 4 & 0 \end{array}$$

Thus, by the remainder theorem, $P(-2) = 0$ and by the factor theorem $x + 2$ is a factor of $P(x)$.

Note Alternatively, we can evaluate

$$\begin{aligned} P(-2) &= (-2)^3 - 3(-2)^2 - 6(-2) + 8 \\ &= -8 - 12 + 12 + 8 \\ &= 0 \end{aligned}$$

2. Find the solution set of the equation $x^3 - 3x^2 - 6x + 8 = 0$.

In example 1, we determined that $x + 2$ is a factor of $x^3 - 3x^2 - 6x + 8$

and the quotient $\frac{P(x)}{x + 2} = x^2 - 5x + 4$.

Thus, we have

$$\begin{aligned} (x + 2)(x^2 - 5x + 4) &= 0 \\ (x + 2)(x - 4)(x - 1) &= 0 \quad \text{Factor the left member} \end{aligned}$$

Since $x = 4$ when $x - 4 = 0$ and $x = 1$ when $x - 1 = 0$, the solution set is $\{-2, 4, 1\}$.

Note We have used the zero product property stated in section 4-1 to find the remaining solutions.

3. Find a polynomial $P(x)$ of degree 3 whose zeros are 2, -1 , and -3 .
By the factor theorem, $(x - 2)$, $(x + 1)$, and $(x + 3)$ are the factors of $P(x)$.
Thus,

$$P(x) = (x - 2)(x + 1)(x + 3) = x^3 + 2x^2 - 5x - 6$$

is a polynomial of degree 3 having the given zeros.

Note This polynomial is *not unique*, since multiplying $P(x)$ by any nonzero real number will yield another polynomial having the same zeros. To illustrate,

$$P(x) = 4x^3 + 8x^2 - 20x - 24$$

also has zeros 2, -1 , and -3 .

► **Quick check** Show that $x + 5$ is a factor of $P(x) = x^3 + 4x^2 - 7x - 10$.
Find the solution set of the equation $x^3 + 4x^2 - 7x - 10 = 0$. ■

Given

$$P(x) = 4x^2 - 12x + 9$$

factoring the polynomial, $P(x) = (2x - 3)(2x - 3)$ and the zeros are $\frac{3}{2}$ and $\frac{3}{2}$. Thus, we may have repeated zeros. In this case, $\frac{3}{2}$ is repeated twice and we say the $P(x)$ has zero $\frac{3}{2}$ of *multiplicity 2*. In general, if a factor $x - r$ is repeated k times in a polynomial, we say r is a *zero of multiplicity k* .

■ Example 4-6 D

- Given $P(x) = (x + 2)^2(x - 5)^3(x - 1)^4$, the distinct zeros are -2 , 5 , and 1 . Since each is repeated more than once, we say -2 is a zero of multiplicity 2, 5 is a zero of multiplicity 3, and 1 is a zero of multiplicity 4.
- Given 3 is a zero of the polynomial $P(x) = 2x^3 - 7x^2 - 7x + 30$, find the other zeros.
Since 3 is a zero, then $x - 3$ is a factor of $P(x)$ by the factor theorem. Using synthetic division,

$$\begin{array}{r|rrrr} 3 & 2 & -7 & -7 & 30 \\ & \downarrow & 6 & -3 & -30 \\ \hline & 2 & -1 & -10 & 0 \end{array}$$

Thus, $P(x) = (x - 3)(2x^2 - x - 10)$ and factoring $2x^2 - x - 10$,

$$P(x) = (x - 3)(2x - 5)(x + 2)$$

Since $x = 5/2$ when $2x - 5 = 0$ and $x = -2$ when $x + 2 = 0$, the zeros of $P(x)$ are 3, $5/2$, and -2 .

3. Given -1 is a solution (root) of the equation

$$3x^4 - 7x^3 - 33x^2 - 33x - 10 = 0$$

of multiplicity 2, find the solution set.

Given -1 is a solution of multiplicity 2, then $(x + 1)^2 = x^2 + 2x + 1$ is a factor of the left member of the equation. We divide

$$\begin{array}{r}
 3x^2 - 13x - 10 \\
 x^2 + 2x + 1 \overline{) 3x^4 - 7x^3 - 33x^2 - 33x - 10} \\
 \underline{3x^4 + 6x^3 + 3x^2} \\
 -13x^3 - 36x^2 - 33x \\
 \underline{-13x^3 - 26x^2 - 13x} \\
 -10x^2 - 20x - 10 \\
 \underline{-10x^2 - 20x - 10} \\
 0
 \end{array}$$

Thus, $(x + 1)^2(3x^2 - 13x - 10) = 0$ and if we factor $3x^2 - 13x - 10$, we have

$$(x + 1)^2(3x + 2)(x - 5) = 0$$

The solution set is $\left\{-1, -\frac{2}{3}, 5\right\}$.

► **Quick check** Given the equation $2x^4 + 5x^3 - 51x^2 + 80x - 28 = 0$ has a solution 2 of multiplicity 2, find the solution set of the equation. Find a polynomial $P(x)$ of lowest degree whose zeros are 3, -2 , and 1 of multiplicity 2.

Mastery points

Can you

- Divide $P(x)$ by $x - r$ using synthetic division?
- Use the remainder theorem to find the remainder R when polynomial $P(x)$ is divided by polynomial divisor $x - r$?
- Use the factor theorem to determine if polynomial $x - r$ is a factor of $P(x)$?
- Use the factor theorem to solve an equation $P(x) = 0$ where $P(x)$ is of degree greater than or equal to 3?
- Use the factor theorem to find an equation when given the real roots of the equation?
- Determine the multiplicity of a factor $x - r$?

Exercise 4-6

Perform the indicated divisions using synthetic division. Assume that no divisor is equal to zero. See example 4-6 A.

Example $(x^3 + 10x - 2) \div (x + 2)$

Solution Since there is no x^2 term, we insert $0x^2$ in the dividend to obtain $(x^3 + 0x^2 + 10x - 2) \div (x + 2)$.

$$\begin{array}{r|rrrr} -2 & 1 & 0 & 10 & -2 \\ & \downarrow & -2 & 4 & -28 \\ \hline & 1 & -2 & 14 & -30 \end{array}$$

← Coefficients
← Add

The answer is $x^2 - 2x + 14 - \frac{30}{x+2}$.

1. $\frac{a^2 + 7a + 10}{a + 5}$

2. $\frac{x^2 + 7x + 6}{x + 1}$

3. $\frac{3a - 10 + a^2}{a - 2}$

4. $\frac{y^2 - 6 + y}{y - 2}$

5. $\frac{x^2 + 5x + 9}{x - 2}$

6. $\frac{y^2 - 6y - 4}{y - 3}$

7. $\frac{2a + 3a^2 - 1}{a + 1}$

8. $\frac{5 - 3x + 2x^2}{x + 2}$

9. $\frac{a^3 + 2a^2 - a - 4}{a - 1}$

10. $\frac{y^3 - 3y^2 + y + 2}{y + 1}$

11. $\frac{x^3 - 1}{x - 1}$

12. $\frac{a^3 + 1}{a + 1}$

13. $\frac{y^3 + 3y^2 - 4}{y - 2}$

14. $\frac{x^3 + 5x^2 - 1}{x - 1}$

15. $\frac{2a^3 - 9a^2 - 8a + 17}{a - 5}$

17. $\frac{x^4 - 5x^3 + 5x^2 + 6x - 10}{x - 3}$

16. $\frac{3x^3 + 11x^2 + 12}{x + 4}$

18. $\frac{2a^4 + 6a^3 + a^2 + 2a - 5}{a + 3}$

19. $\frac{3y^4 - 7y^3 - 10y^2 + 11y + 6}{y - 3}$

20. $\frac{a^5 + 2a^4 + 3a^3 + 6a^2 + a + 2}{a + 2}$

Given $P(x) = 3x^4 - 2x^3 + x^2 - 4x + 1$, in problems 21–26, find the following by (a) using the remainder theorem and (b) by substituting the indicated value of x . See example 4-6 B.

Example Using synthetic division and the remainder theorem, find the remainder when $P(x) = 3x^3 - 2x^2 + 4x - 5$ is divided by $(x + 4)$. Evaluate $P(-4)$ using substitution.

Solution a. Using synthetic division,

$$\begin{array}{r|rrrr} -4 & 3 & -2 & 4 & -5 \\ & & -12 & 56 & -240 \\ \hline & 3 & -14 & 60 & -245 \end{array}$$

← Remainder

b. $P(x) = 3x^3 - 2x^2 + 4x - 5$

$$\begin{aligned} P(-4) &= 3(-4)^3 - 2(-4)^2 + 4(-4) - 5 && \text{Replace } x \text{ with } -4 \\ &= -192 - 32 - 16 - 5 \\ &= -245 \\ P(-4) &= -245 \end{aligned}$$

21. $P(1)$

22. $P(-1)$

23. $P(-3)$

24. $P(3)$

25. $P(0)$

26. $P(4)$

27. Given $P(x) = 5x^5 - 2x^3 + 1$, use the remainder theorem to find $P(2)$.

29. Given $P(x) = 24x^4 - 16x^2 + 14x - 6$, use the remainder theorem to find $P\left(\frac{1}{2}\right)$.

28. Given $P(x) = 8x^6 + 3x^2 - 2x - 5$, use the remainder theorem to find $P(-2)$.

Use the factor theorem to determine if the given binomial is a factor of the polynomial $P(x)$. See example 4-6 C-1.

Example Show that $x + 5$ is a factor of $P(x) = x^3 + 4x^2 - 7x - 10$.

Solution Using synthetic division,

$$\begin{array}{r|rrrr} -5 & 1 & 4 & -7 & -10 \\ & & -5 & 5 & 10 \\ \hline & 1 & -1 & -2 & 0 \end{array} \leftarrow x + 5 \text{ is a factor}$$

30. $P(x) = 3x^2 - 4x - 4; x - 2$

32. $P(x) = x^3 + x^2 - 7x - 10; x + 2$

34. $P(x) = 4x^4 - 13x^3 - 13x^2 - 4x + 12; x + 2$

36. $P(x) = x^4 - 81; x - 3$

38. $P(x) = 3x^5 - 3x^4 + 5x^2 - 13x - 6; x - 3$

31. $P(x) = 2x^2 + 3x - 5; x + 3$

33. $P(x) = x^3 - x^2 + 2x - 8; x - 2$

35. $P(x) = x^4 - 9x^3 + 18x^2 - 3; x + 1$

37. $P(x) = x^3 + 64; x + 4$

39. $P(x) = 3x^5 + 4x^2 - 7; x + 1$

Find the solution set of the following equations using the factor theorem and the given root. See example 4-6 C-2 and D-3.

Example Given the equation $2x^4 + 5x^3 - 51x^2 + 80x - 28 = 0$ has a solution 2 of multiplicity 2, find the solution set of the equation.

Solution Given a solution 2 of multiplicity 2, then $(x - 2)^2 = x^2 - 4x + 4$ is a factor of the equation. We divide $2x^4 + 5x^3 - 51x^2 + 80x - 28$ by $x^2 - 4x + 4$.

$$\begin{array}{r} x^2 - 4x + 4 \overline{) 2x^4 + 5x^3 - 51x^2 + 80x - 28} \\ \underline{2x^4 - 8x^3 + 8x^2} \\ 13x^3 - 59x^2 + 80x \\ \underline{13x^3 - 52x^2 + 52x} \\ -7x^2 + 28x - 28 \\ \underline{-7x^2 + 28x - 28} \\ 0 \end{array}$$

The quotient $2x^2 + 13x - 7 = (2x - 1)(x + 7)$, and if we set each factor equal to zero, we obtain $x = \frac{1}{2}$ and $x = -7$. The solution set is $\left\{-7, 2, \frac{1}{2}\right\}$.

40. $x^3 + 7x^2 - 2x - 12 = 0$; -2

42. $x^3 - 5x^2 - 2x + 24 = 0$; 3

44. $12x^3 + 29x^2 + 8x - 4 = 0$; -2

46. $x^4 + 6x^3 - 3x^2 - 52x - 60 = 0$;
-2 has multiplicity 2

41. $3x^3 + 19x^2 - 38x + 16 = 0$; 1

43. $2x^3 + x^2 - 61x + 30 = 0$; 5

45. $x^4 - 2x^2 + 1 = 0$; 1 has multiplicity 2

Find a polynomial $P(x)$ of lowest degree having the given zeros. See example 4-6 C-3.

Example Find a polynomial $P(x)$ of lowest degree whose zeros are 3, -2, and 1 of multiplicity 2.

Solution Since $P(x)$ has a zero

1. 3, then $x - 3$ is a factor

2. -2, then $x + 2$ is a factor

3. 1 of multiplicity 2, then $(x - 1)^2 = x^2 - 2x + 1$ is a factor

Thus, $P(x) = (x - 3)(x + 2)(x^2 - 2x + 1) = x^4 - 3x^3 - 3x^2 + 11x - 6$.

47. 2, -1, 4

48. 5, -3, 3

49. 4, 0, -1

50. -2, 2, -3, 3

51. -8, 8, -1, 1

52. 4, -3, -7, 7

53. 5, -5, 6, 2

54. 3 of multiplicity 2; -3 of multiplicity 2

55. -1 of multiplicity 2; 4 of multiplicity 2

56. 0 of multiplicity 3; -2, 4

57. -5, -2, 0 of multiplicity 2

In problems 58-61, find the zeros of the given polynomial, indicating multiplicity where necessary. See example 4-6 D-1.

58. $P(x) = (x - 2)^2(x + 3)(x - 4)^2$

59. $P(x) = (x - 2)^2(x + 2)^2(x - 3)^3(x + 4)^2$

60. $P(x) = x^3(x - 3)^2(x + 5)$

61. $P(x) = x^2(x + 4)^4(x - 3)^2(x - 6)$

62. Show that $x - 2$ is a factor of $P(x) = x^{10} - 1024$.

63. Find the remainder when $4x^{98} - 8x^{47} + 9x^{28} - 3x^{17} + 1$ is divided by $x + 1$.

64. Find all values of k such that $P(x) = k^2x^3 - 7kx + 6$ is divisible by $x - 1$.

65. Use synthetic division to decide whether or not $3/2$ is a solution of the equation $2z^4 - 5z^3 + 11z^2 - 14z + 3 = 0$.

66. Use synthetic division to decide whether or not $1/3$ is a solution of the equation $6x^4 + x^3 - 4x^2 + 13x - 4 = 0$.

Review exercises

Find the solution set of the following equations. See section 2-1.

1. $5(x - 4) + 12 = 3x + 2x - 6$

2. $\frac{1}{2}x - 1 = \frac{1}{3}x + 3$

3. $\frac{7}{12}y - 4 = \frac{5}{6}y - 5$

4. One number is 27 more than another number. The lesser number is one-fourth of the greater number. Find the two numbers. See section 2-3.

5. Solve $V = \frac{1}{3}\pi h^2(3R - h)$ for R . See section 2-2.

“Your Company” is helping to reduce college costs...



The Problem: College is expensive. And textbooks rank among the fastest growing costs. According to some sources, students now pay over \$900 a year for course materials. And they are going deeper into debt to afford college. The result is an alarming development: over 50% of today's students are not buying all their required course material, marginalizing their investment in higher education due to textbook costs.

The Solution: By purchasing advertising in the **Freeload Press** suite of publications, commercial and non-profit sponsors deliver their marketing message to today's college students and reduce (or eliminate!) the cost of that textbook for the student.

4-7 ■ Equations containing rational expressions

Rational equations

In chapter 2, we studied how to find the solution set of a linear equation. We now consider equations that contain at least one term that is a rational expression. We call these equations **rational equations**. The same properties of real numbers that we used in chapter 2 will apply to solve rational equations once the polynomial denominators are eliminated.

Recall that an equivalent equation is obtained when each term of an equation is multiplied by the same nonzero number. The following examples demonstrate the use of this property to solve rational equations.

■ **Example 4-7 A**

Find the solution set of the following rational equation.

$$1. \frac{x}{3} + \frac{x}{4} - \frac{x}{6} = 35$$

The LCM of 3, 4, and 6 is 12.

$$\begin{aligned} 12 \cdot \frac{x}{3} + 12 \cdot \frac{x}{4} - 12 \cdot \frac{x}{6} &= 12 \cdot 35 \\ 4x + 3x - 2x &= 420 \\ 5x &= 420 \\ x &= 84 \end{aligned}$$

Multiply each term by the LCM 12
Simplify each term
Combine like terms
Divide by 5

The solution set is $\{84\}$.

When the rational equation contains rational expressions with the variable in the denominator, multiplying all terms by the LCM of the denominators does *not always* produce an equation that is equivalent to the original equation.

$$2. 1 - \frac{2a}{a+1} = \frac{2}{a+1}$$

$$\begin{aligned} (a+1) \cdot 1 - (a+1) \cdot \frac{2a}{a+1} &= (a+1) \cdot \frac{2}{a+1} \\ a+1 - 2a &= 2 \\ 1-a &= 2 \\ -a &= 1 \\ a &= -1 \end{aligned}$$

Multiply each term by the LCM $(a+1)$
Reduce in each term
Combine like terms
Subtract 1 from each member
Multiply each member by -1

However, in the original equation, -1 is *not in the domain of the rational expressions involved*. Therefore -1 cannot be a solution and we conclude that the equation has no solution. Hence the solution set is \emptyset .

► **Quick check** Find the solution set of $3 - \frac{4x}{x-2} = \frac{-8}{x-2}$.

Extraneous solutions

The possible solution -1 in example 2 is called an **extraneous solution**—a solution of the equation obtained by multiplying each term by the LCM of the denominators but *not a solution of the original equation*.

Note It is important to observe the domain of the rational expressions in the rational equation to know where extraneous solutions may occur.

To further illustrate, consider the rational equation

$$\frac{3x}{x-3} = 2 + \frac{9}{x-3}$$

Note that the domain of the rational expressions is all real numbers *except* 3. That is, $x \neq 3$. If we proceed to solve the equation, we *do* obtain $x = 3$. Thus 3 is an extraneous solution, the equation has no solution, and the solution set is \emptyset .

The procedure for solving rational equations is stated here.

To solve rational equations

1. Find the LCM of the denominators.
2. Multiply each term by the LCM of the denominators.
3. Solve the resulting equation.
4. Check for extraneous solutions if the LCM contains a variable.

Equations and formulas containing rational expressions and more than one variable are common in scientific fields. It is often desirable to solve such equations for one variable in terms of the other variables in the equation. We use the same procedure that we used in solving preceding equations of this section.

Example 4-7 B

Solve each of the following equations for the indicated variable. Assume all denominators are nonzero.

1. $\frac{3x}{5} - \frac{4y}{3} = 6$ for x

$$15 \cdot \frac{3x}{5} - 15 \cdot \frac{4y}{3} = 15 \cdot 6$$

$$3 \cdot 3x - 5 \cdot 4y = 90$$

$$9x - 20y = 90$$

$$9x = 90 + 20y$$

$$x = \frac{90 + 20y}{9}$$

Multiply by the LCM of 5 and 3,
namely 15.
Simplify each term.

Add $20y$ to each member.

Divide each member by 9.

2. $\frac{3}{x} - 7 = 9y + \frac{2y}{3x}$ for y

$$3x \cdot \frac{3}{x} - 3x \cdot 7 = 3x \cdot 9y + 3x \cdot \frac{2y}{3x}$$

$$3 \cdot 3 - 21x = 27xy + 2y$$

$$9 - 21x = 27xy + 2y$$

Multiply each term by LCM of
 x and $3x$, $3x$.
Simplify each term.

Since we cannot combine the terms containing y in the right member, we must *factor* the common factor, y , from each term.

$$9 - 21x = (27x + 2)y$$

Factor y in the right member

$$\frac{9 - 21x}{27x + 2} = y$$

Divide each member by the coefficient of y , $(27x + 2)$

3. In physics, the rule governing the speeds of two gears, one the driver gear A and the other the driven gear B , is given by

$$\frac{T_A}{T_B} = \frac{R_B}{R_A}$$

where T = the number of teeth in the gear and R = the revolutions per minute of the gear. Solve for the revolutions per minute of the driven gear, R_B .

$$T_B R_A \cdot \frac{T_A}{T_B} = T_B R_A \cdot \frac{R_B}{R_A}$$

Multiply each term by the LCM of T_B and R_A , $T_B R_A$

$$R_A \cdot T_A = T_B \cdot R_B$$

Reduce in each term

$$R_B = \frac{R_A T_A}{T_B}$$

Divide each member by T_B and interchange the two members

► **Quick check** Given $\frac{6}{p-3} + \frac{2}{pq} = \frac{5}{4p}$, solve for p . ■

Mastery points

Can you

- Solve a rational equation in one variable?
- Solve a rational equation in two or more variables for one of the variables?

Exercise 4-7

Find the solution set of each of the following equations. Where extraneous solutions exist, so state. See example 4-7 A.

Example Find the solution set of $3 - \frac{4x}{x-2} = \frac{-8}{x-2}$.

Solution The LCM is $(x - 2)$.

$$(x-2)3 - (x-2)\frac{4x}{x-2} = (x-2)\frac{-8}{x-2}$$

Multiply each term by LCM $x - 2$

$$3x - 6 - 4x = -8$$

Reduce and multiply

$$-x - 6 = -8$$

Combine like terms

$$-x = -2$$

Add 6 to each member

$$x = 2$$

Multiply by -1

Since the domain of the rational expressions is all real numbers except 2, the solution set is \emptyset and 2 is an extraneous solution.

1. $\frac{x+7}{4} = \frac{2x}{12}$

2. $\frac{4b-3}{10} = \frac{2b-1}{6}$

3. $\frac{3x}{5} - \frac{4x}{3} = 1$

4. $\frac{m}{3} + 4 = \frac{7m}{4}$

7. $\frac{b-3}{10} + \frac{2b+1}{15} = 2$

10. $\frac{5-b}{8b} = \frac{3b+7}{6b}$

13. $\frac{5}{4a+2} = \frac{7}{2a+1}$

16. $1 + \frac{5}{3m-9} = \frac{10}{m-3}$

19. $\frac{x}{x-2} + \frac{2}{3} = \frac{2}{x-2}$

22. $\frac{5y}{3-y} + \frac{8}{y-3} = 4$

25. $\frac{5}{a-3} - \frac{1}{a+2} = \frac{6a}{a^2-a-6}$

27. $\frac{4}{2x-6} - \frac{12}{4x+12} = \frac{12}{x^2-9}$

29. $\frac{6}{q^2+q-6} = \frac{5}{q^2+3q-10}$

31. $\frac{13}{n^2+2n-15} - \frac{1}{n^2+10n+25} = 0$

33. $\frac{4}{x^2-9} = \frac{7}{x^2-7x+12} - \frac{5}{x^2-x-12}$

35. $\frac{6}{2n^2+n-3} - \frac{5}{4n^2-9} = \frac{6}{2n^2-5n+3}$

5. $\frac{4}{a} - \frac{6}{3a} = \frac{3}{5}$

8. $\frac{3y-4}{16} - \frac{2-3y}{12} = 1$

11. $\frac{5}{x-2} = \frac{4}{2x+1}$

14. $\frac{11}{3y-2} = \frac{8}{12y-8}$

17. $\frac{x-1}{x^2-4} = \frac{6}{x-2}$

20. $\frac{3}{2} - \frac{1}{x-4} = \frac{-2}{2x-8}$

23. $\frac{8b}{b^2-16} = \frac{3}{b+4} + \frac{5}{4-b}$

6. $4 - \frac{5}{9b} = \frac{7}{6b}$

9. $\frac{x-3}{3x} = \frac{2x+3}{9x}$

12. $\frac{-3}{x+7} = \frac{2}{3x-1}$

15. $\frac{6}{6x+3} = \frac{3}{2x+1} + 5$

18. $\frac{5}{y+3} = \frac{2y+1}{y^2-9}$

21. $4 - \frac{2x}{5-x} = \frac{6}{x-5}$

24. $\frac{10}{5-a} = \frac{7}{a+5} - \frac{6}{a^2-25}$

26. $\frac{7}{x+6} = \frac{9}{x-9} - \frac{2x-1}{x^2-3x-54}$

28. $\frac{11x-3}{10x^2-3x-4} - \frac{4}{5x-4} = \frac{9}{2x+1}$

30. $\frac{9}{y^2-6y+8} = \frac{14}{y^2-16}$

32. $\frac{2}{2n^2-7n-4} + \frac{6}{6n^2-5n-4} = 0$

34. $\frac{-2}{a^2+3a-4} - \frac{6}{a^2-1} = \frac{1}{a^2+5a+4}$

36. $\frac{6}{8x^2-14x-15} + \frac{1}{8x^2-26x+15} = \frac{9}{16x^2-9}$

Solve the given equations and formulas for the indicated variable. Assume all denominators are nonzero. See example 4-7 B.

Example Given $\frac{6}{p-3} + \frac{2}{pq} = \frac{5}{4p}$, solve for p .

Solution The LCM of $4p$, pq , and $p-3$ is $4pq(p-3)$.

$$4pq(p-3) \cdot \frac{6}{p-3} + 4pq(p-3) \cdot \frac{2}{pq} = 4pq(p-3) \cdot \frac{5}{4p}$$

$$4pq \cdot 6 + 4(p-3)2 = q(p-3)5$$

$$24pq + 8(p-3) = 5q(p-3)$$

$$24pq + 8p - 24 = 5pq - 15q$$

$$19pq + 8p = 24 - 15q$$

$$(19q + 8)p = 24 - 15q$$

$$p = \frac{24 - 15q}{19q + 8}$$

$$p = \frac{3(8 - 5q)}{19q + 8}$$

Multiply each term by LCM $4pq(p-3)$.

Reduce in each term.

Commute and multiply.

Distributive property.

Add 24 to and subtract $5pq$ from each member.

Factor p in right member.

Divide each member by $19q + 8$.

Factor numerator (not reducing).

37. $\frac{4}{a} - \frac{3}{b} = 7$ for b

38. $\frac{6}{x} - 4 = \frac{7}{y}$ for y

39. $\frac{3}{p} + 4 = \frac{6q}{2p} - 3a$ for p

40. $\frac{a+6}{3} - \frac{b-2}{4} = \frac{c}{12}$ for a

41. $\frac{y-3}{x+2} = \frac{5}{3}$ for y

42. $\frac{-7}{3} = \frac{y+1}{x-5}$ for x

43. $S = \frac{a}{1-r}$ for r (progression formula)

44. $S = \frac{n(a+b)}{2}$ for a (progression formula)

45. $A = \frac{1}{2}h(b_1 + b_2)$ for b_1 (area of a trapezoid)

46. $C = \frac{5}{9}(F - 32)$ for F (temperature conversion)

See example 4-7 B-3.

47. The coefficient of linear expansion, k , of a solid when heated is given by

$$k = \frac{L_t - L_0}{L_0 t}$$

where L_t is the length at $t^\circ\text{C}$, L_0 is the length at 0°C , and t is any given temperature in Celsius. Solve for t . Solve for L_0 .

48. A formula for unknown resistance R_x in a battery is given by

$$R_x = R_m \left(\frac{E_1}{E_2} - 1 \right)$$

Solve for E_1 .

49. The interest rate, r , on a given amount of money, P , over a given period of time, t , which pays interest, I , is given by

$$r = \frac{I}{Pt}$$

Solve the equation for t .

50. The kinetic energy of a body, KE , is computed by

$$KE = \frac{Wv^2}{2g}$$

where W is the weight in pounds, v is the velocity expressed in feet per second, and g is the acceleration due to gravity. Solve for g .

51. The total resistance R of a parallel circuit is given by

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$$

where R_1 , R_2 , and R_3 are the resistances of the respective circuits. Solve for R_3 .

52. The capacitance of capacitors C_1 , C_2 , and C_3 in a series circuit is given by

$$\frac{1}{C} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3}$$

Solve for C .

53. Given a large piston and a small piston on which forces F and f , respectively, are applied, then

$$\frac{F}{f} = \frac{A}{a}$$

where A is the area of the large piston and a is the area of the small piston. Solve for A .

54. As gas expands when heated, the relationship between pressure P , volume V , and absolute temperature T is given by

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

called Charles' Law. Solve for V_2 .

55. The formula

$$\frac{R_2}{R_1} = \frac{M + T_2}{M + T_1}$$

gives a relationship for the increase in the resistance of a circuit caused by a rise in temperature. Solve for T_2 . Solve for R_1 .

56. The lens maker's formula for the focal length, f , of a lens is given by

$$\frac{1}{f} = (\mu_m - 1) \left(\frac{1}{R_1} + \frac{1}{R_2} \right)$$

where μ_m is the index of refraction and R_1 and R_2 are radii of curvature. Solve for f . Solve for μ_m .

57. If Ann can paint the house in 10 hours, write a rational expression for what part of the house she can paint in (a) 1 hour, (b) 5 hours, (c) t hours.

Learning the hard way?

www.FreeLoadPress.com

HARD

EASY

How will you learn those terms and definitions? Still trying the **hard** way? Memory studies prove that you will forget **80%** of what you learned that way.

Books are a great reference. But when it comes to learning those important terms, why not make it **easy** on yourself with **exciting, multimedia study guides** from Total Recall Learning, where you will truly **remember everything** you learn!

Download the latest Total Recall study guides from www.FreeLoadPress.com to ensure your success in school or college. Getting an "A" has never been so **easy** and so much **fun**!

Information is broken down into **small**, easy-to-digest **pieces**. You will never want to learn any other way again! And with **Know-it-All Live!** you can even prepare your very own multimedia flashcards! For an even greater selection of courses visit www.TotalRecallLearning.com. Make learning **easy**!



www.TotalRecallLearning.com

58. If John can bake a batch of cookies in 3 hours, write a rational expression for what part of the batch he can bake in (a) 1 hour, (b) t hours, (c) 20 minutes.
59. An automobile travels 200 miles. Write a rational expression for the speed (rate of travel) the automobile is traveling if the distance is covered in (a) 5 hours, (b) 4 hours, (c) t hours. (*Hint:* distance = rate \cdot time)
60. Write a rational expression for the time t a boat travels 20 miles downstream if the boat is traveling downstream at (a) 4 mph, (b) 12 mph, (c) r mph.
61. If one number is three times as large as a second number, and if n represents the second number, (a) write a rational expression for the reciprocal of the two numbers, (b) write a rational expression for five times the reciprocal of the second number.
62. If a number n is added to the numerator and the denominator of the rational number $\frac{3}{5}$, write a rational expression for the new number.
63. If the same number n is added to the numerator and taken away from the denominator of $\frac{5}{6}$, write a rational expression for the new number.

Review exercises

Completely factor the following expressions. See sections 3-5, 3-6, and 3-7.

1. $4p^2 - 25q^2$
2. $x^2 - 24x + 144$
3. $2y^2 - 15y - 8$
4. Dene invests \$20,000, part at 6% and the rest at 8% interest. If the total income from the investments in one year is \$1,360, how much did she invest at each rate? See section 2-3.
5. Five times a number increased by 13 gives 53. Find the number. See section 2-3.
6. Given polynomial $P(x) = 2x^2 + x - 3$, find $P(-3)$. See section 1-5.

4-8 ■ Problem solving with rational equations

Now that we can solve rational equations, let us use that skill to solve some types of word problems that result in rational equations when we use problem-solving techniques.

■ Example 4-8 A

Work problem

1. Jan can produce a part in 3 hours and Joe can produce the same part in 4 hours. How long would it take them to produce the part working together? Since Jan can produce the part in 3 hours and Joe can produce the part in 4 hours, then Jan can produce $\frac{1}{3}$ of the part in 1 hour and Joe can produce $\frac{1}{4}$ of the part in 1 hour.
- Let x = the time in hours necessary to produce the part working together.

Then they can produce $\frac{1}{x}$ of the part in 1 hour.

$$\begin{array}{ccccc} \text{Jan in 1 hour} & + & \text{Joe in 1 hour} & = & \text{together in 1 hour} \\ \frac{1}{3} & + & \frac{1}{4} & = & \frac{1}{x} \end{array}$$

The LCM of the denominators is $12x$.

$$\begin{aligned} 12x \cdot \frac{1}{3} + 12x \cdot \frac{1}{4} &= 12x \cdot \frac{1}{x} && \text{Multiply each term by the LCM } 12x. \\ 4x + 3x &= 12 && \text{Reduce in each term.} \\ 7x &= 12 && \text{Add in left member.} \\ x &= \frac{12}{7} \text{ or } 1\frac{5}{7} \text{ hours} && \text{Divide by 7.} \end{aligned}$$

Working together, Jan and Joe can produce the part in $1\frac{5}{7}$ hours.

Uniform motion problems

2. An automobile can travel 200 miles in the same time that a truck can travel 150 miles. If the automobile travels at an average rate of 15 miles per hour faster than the truck, find the average rate of each. We use the relationship between distance traveled, rate of travel, and time traveled—distance (d) = rate (r) · time (t).

The following table will be helpful in finding the equation necessary to solve this type of problem.

	d	r	t
Automobile			
Truck			

Let r = the rate of the truck. Then $r + 15$ = the rate of the automobile.

	d	r	t
Automobile	200	$r + 15$	$\frac{200}{r + 15}$
Truck	150	r	$\frac{150}{r}$

Times are equal

Note To find the time, we used the relationship

$$\text{time} = \frac{\text{distance}}{\text{rate}}$$

Since both vehicles traveled the same length of time, we obtain the equation

$$\frac{200}{r+15} = \frac{150}{r}$$

$$r(r+15) \cdot \frac{200}{r+15} = r(r+15) \cdot \frac{150}{r}$$

Multiply each term by the LCM of r and $(r+15)$, $r(r+15)$.

$$200r = (r+15)150$$

$$200r = 150r + 2,250$$

$$50r = 2,250$$

$$r = 45$$

$$r+15 = 60$$

Thus the average rate of the truck is 45 miles per hour and of the automobile is 60 miles per hour.

3. John's boat travels at 20 miles per hour in still water. If the speed of the current is 5 miles per hour, how far can John travel downstream if it takes him a total time of 3 hours to go downstream and back upstream?

Let x = the distance traveled downstream (and back upstream since they are the same distance). Using $t = \frac{d}{r}$, we summarize the information in the table.

	d	r	t
Downstream	x	$20 + 5 = 25$	$\frac{x}{25}$
Upstream	x	$20 - 5 = 15$	$\frac{x}{15}$

Total time is 3 hours

Note When traveling downstream, we *add* the speed in still water to the speed of the current. When traveling upstream, we *subtract* the speed of the current from the speed in still water.

Since total time downstream and back upstream is 3 hours, the equation is

$$\frac{x}{25} + \frac{x}{15} = 3 \quad \leftarrow \text{Total time}$$

Downstream \uparrow \uparrow Upstream

$$\frac{x}{25} \cdot 75 + \frac{x}{15} \cdot 75 = 3 \cdot 75$$

Multiply each term by the LCM of 25 and 15, 75.

$$3x + 5x = 225$$

Multiply as indicated.

$$8x = 225$$

Combine like terms.

$$x = \frac{225}{8}$$

Divide each member by 8.

$$= 28\frac{1}{8}$$

John can travel $28\frac{1}{8}$ miles downstream.

Number and reciprocal problem

4. One number is twice another number. The sum of their reciprocals is $\frac{9}{2}$. Find the numbers.

Let x = one number, then $2x$ = the other number. Their reciprocals are then $\frac{1}{x}$ and $\frac{1}{2x}$. The sum of the reciprocals is $\frac{9}{2}$, so we have the equation

$$\frac{1}{x} + \frac{1}{2x} = \frac{9}{2}$$

$$2x \cdot \frac{1}{x} + 2x \cdot \frac{1}{2x} = 2x \cdot \frac{9}{2} \quad \text{Multiply each term by the LCM of } x, 2, \text{ and } 2x, 2x$$

$$2 + 1 = 9x$$

$$3 = 9x$$

$$x = \frac{3}{9} = \frac{1}{3}$$

$$2x = \frac{2}{3}$$

The two numbers are $\frac{1}{3}$ and $\frac{2}{3}$.

Check: Since the sum of the reciprocals $3 + \frac{3}{2} = \frac{6}{2} + \frac{3}{2} = \frac{9}{2}$, the answer is correct.

► **Quick check** Pam can clean her house in 2 hours. Her daughters, Linnea and Jean, can do the job in 2 hours and 3 hours, respectively. At these rates, how long would it take the three of them to clean the house working together?

Mastery points

Can you

- Set up and solve work problems?
- Set up and solve uniform motion problems?
- Set up and solve number and reciprocal problems?

Exercise 4-8

Solve the following work problems. See example 4-8 A-1.

Work problems

Example Pam can clean her house in 2 hours. Her daughters, Linnea and Jean, can do the job in 2 hours and 3 hours, respectively. At these rates, how long would it take the three of them to clean the house working together?

Solution Let x = time in hours to clean the house when the three of them are working together.

Then 1. Pam can clean $\frac{1}{2}$ of the house in 1 hour.

2. Linnea can clean $\frac{1}{2}$ of the house in 1 hour.

3. Jean can clean $\frac{1}{3}$ of the house in 1 hour.

4. Working together, they can clean $\frac{1}{x}$ of the house in 1 hour.

$$\begin{array}{ccccccc} \text{part done} & & \text{part done} & & \text{part done} & & \text{part done by all} \\ \text{by Pam} & & \text{by Linnea} & & \text{by Jean} & & \text{three working together} \\ \frac{1}{2} & + & \frac{1}{2} & + & \frac{1}{3} & = & \frac{1}{x} \end{array}$$

The equation is $\frac{1}{2} + \frac{1}{2} + \frac{1}{3} = \frac{1}{x}$.

$$6x \cdot \frac{1}{2} + 6x \cdot \frac{1}{2} + 6x \cdot \frac{1}{3} = 6x \cdot \frac{1}{x}$$

$$3x + 3x + 2x = 6$$

$$8x = 6$$

$$x = \frac{6}{8}$$

$$x = \frac{3}{4}$$

Multiply each term by the LCM of 2, 3, and x , $6x$.

Combine like terms.

Divide each member by 8.

Simplify.

It would take Pam and her daughters $\frac{3}{4}$ of an hour to clean the house working together.

- Pete Hansen milks his herd of cattle in 2 hours, whereas his son John milks them in 3 hours. How many hours and minutes would it take them to do the same job working together?
- Jake has two hay balers, A and B . If Jake can completely bale a given field in 3 hours using baler A and in $3\frac{3}{4}$ hours using baler B , how long would it take him to bale the field working both balers together?
- During a "clean-up, paint-up, fix-up week" in Sault Sainte Marie, Jim, Toni, and Ken clean up a given vacant lot. If it takes each person $1\frac{1}{2}$ hours, 2 hours, and 3 hours, respectively, to do the job alone, how long would it take them working together to clean up the lot?
- Tanya, Bill, and Neysa work in a bakery and can mix 12 loaves of bread individually in 36 minutes, 40 minutes, and 30 minutes, respectively. If they worked together, in how many minutes could they mix the 12 loaves?
- If two water pipes can fill a swimming pool in 18 hours when both pipes are open and one of the pipes can fill the pool alone in 30 hours, how long would it take the other pipe to fill the pool alone?
- In a machine shop, machine A can produce certain parts in 1 hour and 20 minutes. If machines A and B working together can produce the parts in 50 minutes, how long would it take machine B to do the job alone?
- Two inlet pipes can fill a water basin in 10 hours and 12 hours, respectively, when open individually. If an outlet pipe can empty the basin alone in 9 hours, how long would it take to fill the basin if all three pipes are open simultaneously?
- An open sink drain can empty a sink full of water in 2 minutes. If the cold water and hot water faucets can, when open fully, fill the sink in $3\frac{1}{2}$ and 3 minutes, respectively, how long would it take to fill the sink if all three are open simultaneously?
- It takes pump A twice as long to unload an oil tanker as it takes pump B to do the job. If the two pumps working together can unload the tanker in 12 hours, how long will it take each to do the job?

10. If one microprocessor can process a set of inputs in three-fifths of the time that a second microprocessor can do the job, and together the machines can do the job in 2 milliseconds, how long will it take each microprocessor to process the set of inputs individually?
11. It takes tug *A* one-half as long to push a series of barges up a river as it takes tug *B* to do the same job. If it takes tug *C* two times as long to do the job as it does tug *B* and all three tugs working together can do the job in 4 hours, how long would it take tug *A* to do the job alone?
12. A portable gasoline-powered generator, a solar cell, and a wind generator fully charge a dead storage battery in 9 hours when charging simultaneously. If the wind generator takes twice as long to fully charge the battery as the solar cell takes and the gasoline-powered generator takes three-fourths as long as the solar cell, how long would it take the solar cell to fully charge the battery alone?

Solve the given uniform motion problems. See example 4-8 A-2 and 3.

Uniform motion problems

13. An excursion boat moves at 16 miles per hour in still water. If the boat travels 20 miles downstream in the same time it takes to travel 14 miles upstream, what is the speed of the current? (*Hint:* Let x = the speed of the current. Then $16 + x$ = the speed of the boat downstream and $16 - x$ = the speed of the boat upstream.)
14. An airplane can cruise at 300 miles per hour in still air. If the airplane takes the same time to fly 950 miles with the wind as it does to fly 650 miles against the wind, what is the speed of the wind?
15. A boat travels 40 kilometers upstream in the same time that it takes the same boat to travel 60 kilometers downstream. If the stream is flowing at 6 kilometers per hour, what is the speed of the boat in still water?
16. The speed of a wind is 15 miles per hour. Find the speed of an airplane in still air if it flies 200 miles against the wind in the same time that it flies 300 miles with the wind.
17. On a trip from Detroit to Los Angeles, a TWA plane takes one-third of an hour longer to make the trip than an American Airlines plane does. If the TWA plane flies at 300 miles per hour and the American Airlines plane flies at 320 miles per hour, how far is it from Detroit to Los Angeles?
18. It takes Bonnie 2 minutes longer to jog a certain distance than it does Maryann. What distance did they run if Bonnie can jog at 5 miles per hour and Maryann can jog at 7 miles per hour?
19. Mary can row her boat 3 miles per hour in still water. If the current of the river is 2 miles per hour, how far downstream can she row if it takes her 2 hours to go down and back?
20. Peter can average 10 miles per hour riding his bike to deliver his papers. By car he can average 30 miles per hour. If it takes him $\frac{1}{2}$ hour less time by car, how long is Peter's paper route?

Solve the following number and reciprocal problems. See example 4-8 A-4.

Number and reciprocal problems

21. One number is three times another number. The sum of their reciprocals is 2. Find the numbers.
22. One number is twice another number. The sum of their reciprocals is $\frac{15}{8}$. Find the numbers.
23. If the same number is added to the numerator and the denominator of the fraction $\frac{3}{7}$, the result is $\frac{4}{5}$. Find the number.
24. If the same number is added to the numerator and the denominator of the fraction $\frac{1}{4}$, the result is $\frac{2}{3}$. Find the number.
25. What number must be added to the denominator of $\frac{6}{7}$ to obtain $\frac{3}{5}$?
26. What number must be added to the numerator and subtracted from the denominator of $\frac{5}{8}$ to obtain its reciprocal?

3. On the Menu Bar, click Bookmarks.



4. When this menu appears, click Google.



Show is better than Tell.

Visibooks computer textbooks use large screenshots and very little text. They work great for beginners, and people who want to learn computer subjects quickly.

Visibooks are used in college computer classes from Boston College to UC San Diego.

To learn more about them, visit www.visibooks.com.

Page from a typical Visibook:
12 words, two large pictures.



The following word problems are a mixture of the previous types.

27. A boat can go 6 miles upstream in the same time as it can go 10 miles downstream. If the boat goes 12 miles per hour in still water, what is the speed of the current?
28. What number must be added to the numerator of the fraction $\frac{11}{15}$ to obtain the fraction $\frac{7}{5}$?
29. Jane can type an English term paper in 3 hours. If Pamela helps her, together they can type the paper in 2 hours. How long would it take Pamela to type the paper alone?
30. A water tank can be filled by its inlet pipe in 45 minutes. If its outlet pipe can empty the tank in 30 minutes, how long would it take to empty a full tank if both pipes are open?
31. Fran traveled 120 miles to visit her aunt. If she drove twice as fast returning home and the return trip took 2 hours less time, what was her speed going to her aunt's?
32. If one number is three times a second number and the sum of their reciprocals is $\frac{2}{9}$, find the numbers.
33. Erin can jog 2 miles per hour faster than her cousin Sarah. Erin can jog 9 miles in the same time Sarah can jog 6 miles. How fast does each girl jog?

Review exercises

Perform the indicated operations. See section 3-2.

1. $4x(2x^2 - 3x + 1)$ 2. $(3x - 1)(x - 5)$ 3. $(5z - 4)^2$ 4. $(2y + 3)(2y - 3)$

Simplify the following expressions. Express the answer with positive exponents. See section 3-3.

5. $y^{-3} \cdot y^2 \cdot y^0$ 6. $\frac{x^3y^2}{x^{-1}y^{-1}}$
7. What number(s) when squared yield (yields) 49? See section 3-1.
8. What number cubed yields -125? See section 3-1.

Chapter 4 lead-in problem

Doug Nance opens the valve to fill his new swimming pool. Without his knowledge, the drain to empty the pool has been left open. If the pool would normally fill in 10 hours and the drain can empty the full pool in 15 hours, how long does it take to fill the pool?

Solution

Let t represent the time to fill the pool. In 1 hour,

- a. the valve would fill $\frac{1}{10}$ of the pool.
- b. the drain would empty $\frac{1}{15}$ of the pool.
- c. with both open, $\frac{1}{t}$ of the pool would be filled.

We subtract the drain time from the fill time to get the equation

$$\begin{aligned} \frac{1}{10} - \frac{1}{15} &= \frac{1}{t} \\ 30t\left(\frac{1}{10} - \frac{1}{15}\right) &= 30t \cdot \frac{1}{t} && \text{Multiply each member by the LCD } 30t \\ 30t\left(\frac{1}{10}\right) - 30t\left(\frac{1}{15}\right) &= 30 && \text{Distribute in left member and multiply} \\ 3t - 2t &= 30 && \text{Simplify in left member} \\ t &= 30 && \text{Combine like terms} \end{aligned}$$

It takes 30 hours to fill the pool.

Chapter 4 summary

1. A **rational expression** is any algebraic expression that can be written as a quotient of two polynomials where the denominator does not equal zero.
2. The **domain** of a rational expression is the set of all replacement values of the variable for which the expression is defined.
3. The **fundamental principle of rational expressions** states that we obtain an equivalent expression when we multiply or divide the numerator and the denominator of a rational expression by the same nonzero polynomial.
4. A rational expression is **reduced to its lowest terms** when the numerator and the denominator have no common polynomial factors other than 1 or -1 .

5. To **multiply** two or more rational expressions, multiply the numerators to determine the numerator of the product and multiply the denominators to determine the denominator of the product.

6. Given polynomials P , Q , R , and S ,

$$\frac{P}{Q} \div \frac{R}{S} = \frac{P}{Q} \cdot \frac{S}{R} = \frac{P \cdot S}{Q \cdot R}$$

where $Q \neq 0$, $R \neq 0$, and $S \neq 0$.

7. Given polynomials P , Q , and R , $Q \neq 0$,

$$\frac{P}{Q} + \frac{R}{Q} = \frac{P + R}{Q}$$

and

$$\frac{P}{Q} - \frac{R}{Q} = \frac{P - R}{Q}$$

8. Given polynomials P , Q , R , and S , $Q \neq 0$ and $S \neq 0$,

$$\frac{P}{Q} + \frac{R}{S} = \frac{P \cdot S + Q \cdot R}{Q \cdot S}$$

and

$$\frac{P}{Q} - \frac{R}{S} = \frac{P \cdot S - Q \cdot R}{Q \cdot S}$$

9. To find the least common multiple (LCM) of a set of polynomials,
 - a. completely factor the polynomials,
 - b. the LCM consists of the product of all distinct factors of the polynomials, each raised to the greatest power to which it appears in any of the factorizations.

10. Rational expressions can be added or subtracted by building each expression to an equivalent rational expression such that they have a common denominator—the least common denominator (LCD).
11. A **complex rational expression** is a rational expression whose numerator or denominator, or both, contain rational expressions.

12. In the complex rational expression

$$\frac{\frac{4}{5}}{\frac{7}{8}}$$

we call $\frac{4}{5}$ the **primary numerator**, $\frac{7}{8}$ the **primary denominator**, and 5 and 8 the **secondary denominators**.

13. To simplify a complex rational expression, multiply the primary numerator and the primary denominator by the least common multiple (LCM) of the secondary denominators.
14. To **divide** a polynomial by a monomial, simply divide each term of the polynomial by the monomial and add or subtract the resulting quotients.
15. When the divisor is a binomial of the form $x - k$, we can use synthetic division to divide a polynomial by a polynomial.
16. The **remainder theorem** and the **factor theorem** uses synthetic division to find (a) the remainder when a polynomial is divided by $x - k$ and (b) the solution set of a polynomial equation.
17. A **rational equation** is an equation that contains at least one expression of the form $\frac{P}{Q}$, $Q \neq 0$, where P and Q are polynomials.
18. To obtain a simple equation, an equation containing no rational expression, multiply both members of the rational equation by the LCM of the denominators.

Chapter 4 error analysis

1. Placing restrictions on the variable in a rational expression

Example: What restrictions must be placed on the

variable in $\frac{3y+2}{4y^2+3y}$?

Since $4y^2 + 3y = 0$ when $y = 0$, then $y \neq 0$.

Correct answer: $y \neq 0$ or $y \neq -\frac{3}{4}$

What error was made? (see page 155)

2. Finding the domain of a rational expression

Example: $\frac{2x-1}{2x^2+5x-3} = \frac{2x-1}{(2x-1)(x+3)} = \frac{1}{x+3}$

Domain = $\{x | x \in R, x \neq -3\}$

Correct answer: Domain = $\{x | x \in R, x \neq -3, \frac{1}{2}\}$

What error was made? (see page 155)

3. Subtracting rational expressions

Example: $\frac{4y+1}{y^2+3} - \frac{2y-4}{y^2+3}$

$= \frac{4y+1-2y-4}{y^2+3} = \frac{2y-3}{y^2+3}$

Correct answer: $\frac{2y+5}{y^2+3}$

What error was made? (see page 167)

4. Adding rational expressions

Example: $\frac{2x}{x-2} + \frac{x-7}{2-x} = \frac{2x+x-7}{x-2} = \frac{3x-7}{x-2}$

Correct answer: $\frac{x+7}{x-2}$

What error was made? (see page 171)

5. Simplifying complex rational expression to lowest terms

Example: Simplify $\frac{1 + \frac{1}{x}}{x - \frac{1}{x}}$.

$\frac{\left(1 + \frac{1}{x}\right) \cdot x}{\left(x - \frac{1}{x}\right) \cdot x} = \frac{x+1}{x^2-1}$

Correct answer: $\frac{1}{x-1}$

What error was made? (see page 176)

6. Reducing a rational expression to lowest terms

Example: $\frac{2y^3+y}{y} = \frac{2y^3+y}{y} = \frac{2y^3+1}{1} = 2y^3+1$

Correct answer: $2y^2+1$

What error was made? (see page 157)

7. Dividing a polynomial by a polynomial

Example: $(3x^3 + 2x^2 - x + 1) \div (x - 1)$

$$\begin{array}{r} x^2 + x - 2 \\ x-1 \overline{) 3x^3 + 2x^2 - x + 1} \\ \underline{x^3 - x^2} \\ 2x^2 - x \\ \underline{x^2 - x} \\ -2x + 1 \\ \underline{-2x + 2} \\ 3 \end{array}$$

Correct answer: $x^2 + 3x + 2 + \frac{3}{x-1}$

What error was made? (see page 185)

8. Solving rational equations

Example: Find the solution set of $\frac{3x}{x-4} = \frac{12}{x-4} + 1$.

$(x-4) \cdot \frac{3x}{x-4} = (x-4) \cdot \frac{12}{x-4} + (x-4) \cdot 1$

$3x = 12 + x - 4$

$2x = 8$

$x = 4 \quad \{4\}$

Correct answer: The solution set is \emptyset .

What error was made? (see page 199)

9. Using synthetic division

Example: Divide $3x^3 - 2x^2 + 4x - 1$ by $x + 3$.

$$\begin{array}{r} 3 \overline{) 3 \quad -2 \quad 4 \quad -1} \\ \underline{9 \quad 21 \quad 75} \\ 3 \quad 7 \quad 25 \quad 74 \end{array} = 3x^2 + 7x + 25 + \frac{74}{x+3}$$

Correct answer: $3x^2 - 11x + 37 - \frac{112}{x+3}$

What error was made? (see page 190)

10. Finding the remainder using synthetic division

Example: Given $P(x) = 4x^2 - 2x + 1$, find $P(3)$.

$$\begin{array}{r} 3 \overline{) 4 \quad -2 \quad 1} \\ \underline{12 \quad -42} \\ 4 \quad -14 \quad 43 \end{array} \quad P(3) = 43$$

Correct answer: $P(3) = 31$

What error was made? (see page 191)

Chapter 4 critical thinking

If n is an integer, for what values of n will $4n^2 + 4n - 3$ represent an odd integer?

Chapter 4 review

[4-1]

State the domain of the given rational expression in set-builder notation.

1. $\frac{5}{x+7}$

2. $\frac{2x+1}{3x-4}$

3. $\frac{4-2x}{x^2-10x+25}$

4. $\frac{4a^2+1}{9a^2-16}$

5. $\frac{3z-1}{6z^2+13z-5}$

6. $\frac{4y-11}{9y^2-12y+4}$

Reduce each rational expression to lowest terms. Assume all denominators are nonzero.

7. $\frac{a^4b^4c}{a^2b^2c^3}$

8. $\frac{-10m^3n^3p}{35m^4np^5}$

9. $\frac{5x-10}{6x-12}$

10. $\frac{24a}{8a^2-16a}$

11. $\frac{5x-10y}{4y^2-x^2}$

12. $\frac{y^3-64}{y^2-16}$

13. $\frac{a^2-24a+144}{a^2-11a-12}$

14. $\frac{4x^2-5x-6}{5x^2-11x+2}$

15. $\frac{10y^2+9y-9}{12-2y-30y^2}$

[4-2]

Perform the indicated multiplication or division. Place necessary restrictions on the variables.

16. $\frac{14x}{3y} \cdot \frac{9y^2}{7x^2}$

17. $\frac{16a^2}{7y} \div \frac{4a}{21y^2}$

18. $\frac{p^2-16}{4p-3} \cdot \frac{16p^2-9}{3p+12}$

19. $\frac{z^2+1}{2z-6} \div \frac{z^4-1}{z^2-6z+9}$

20. $\frac{m^3-8}{m^2-3m} \cdot \frac{m^2+3m-18}{m^2+3m-10}$

21. $\frac{5a^2+17a+6}{10a^2-a-2} \cdot \frac{2a^2+13a-7}{a^2+10a+21}$

22. $\frac{x^2-49}{8x^3+1} \div \frac{x^2-5x-14}{4x^2-1}$

23. $\frac{4x^3-5x^2}{4x+5} \div (16x^2-25)$

24. $\frac{7y^2+10y-8}{y^2+4y+4} \cdot \frac{y^2+6y+9}{7y^2+17y-12}$

25. $\frac{mq-mp-nq+np}{2mq-2mp-nq+np} \div \frac{qm+qn+pm+pn}{2mp-np+2qm-nq}$

[4-3]

Find the least common multiple (LCM) of the set of polynomials.

26. $15xy^2, 20x^3y, 9x^2y^3$

27. $2x+4, 6x^2, x^2-16$

28. $a^2+3a-10, a^2-2a, 3a+15$

29. $p^2-25, p^2+10p+25, p^2-5p$

Perform the indicated additions and/or subtractions. Assume all denominators are nonzero.

30. $\frac{3x}{4y} + \frac{8x}{3y}$

31. $\frac{n-7}{n+4} - \frac{3n+8}{n-1}$

32. $\frac{10}{p^2+7p-18} + \frac{9}{p^3-4p}$

33. $(2b+7) - \frac{b+9}{3b-2}$

34. $\frac{3y-4}{y^2-49} + \frac{2y+1}{y-7}$

35. $\frac{2x}{2x^2-4x-48} - \frac{5x+1}{x^2-36}$

36. $\frac{7}{2a-4} + \frac{9}{a-2} - \frac{6}{2-a}$

37. $\frac{x+2}{x^2-3x-28} - \frac{3x-7}{4x+16} - \frac{5-x}{8x-56}$

38. $\frac{a+b}{a^2-ab-2b^2} + \frac{3a-b}{a^2-4b^2}$

39. In electricity, the total resistance R_t of any parallel circuit is given by

$$\frac{1}{R_t} = \frac{1}{E_1} + \frac{1}{E_2} + \frac{1}{E_3}$$

Combine the expression in the right member.

[4-4]

Simplify each complex rational expression. Assume all denominators are nonzero.

40. $\frac{\frac{4}{a^2}}{\frac{2}{a}}$

41. $\frac{\frac{5}{x-3}}{\frac{4}{x}}$

42. $\frac{\frac{3x}{x-5}}{\frac{x}{x+2}}$

43. $\frac{7 - \frac{3}{b-5}}{8 + \frac{6}{b-5}}$

44. $\frac{\frac{x-y}{2} + \frac{3}{x}}{\frac{3}{y}}$

45. $\frac{(p-3) + \frac{2}{p-1}}{(p+1) - \frac{3}{p-1}}$

[4-5]

Perform the indicated divisions. Assume that no divisor is equal to zero.

46. $\frac{25a^7 + 15a^3 + 10a}{5a}$

47. $\frac{36a^5b^5c^3 - 18a^2b^3c^4 + 6a^2b^3c}{6a^2b^3c}$

48. $(3x^3 + x + 4) \div (x + 1)$

49. $\frac{2x^4 + x^3 - 3x^2 + 3x - 1}{2x - 1}$

[4-6]

Use synthetic division and the remainder theorem to evaluate each polynomial for the given value of the variable.

50. $P(x) = x^3 - 2x^2 + 8x - 3; x = -2$

51. $P(y) = 4y^3 + 2y^2 - y + 6; y = 1$

52. $P(z) = -3z^4 - z^2 + 5z - 3; z = -1$

Use synthetic division to determine if the given number is a solution of the equation.

53. $x^3 + 2x^2 + 3x + 18 = 0; -3$

54. $2y^3 + 4y^2 + y - 3 = 0; -1$

55. $4z^3 - 10z^2 - z - 6 = 0; 2$

[4-7]

Find the solution set of each of the following rational equations.

56. $6 - \frac{10}{9a} = \frac{5}{12a}$

57. $\frac{4m-3}{15} + \frac{2-5m}{20} = 1$

58. $\frac{4}{a+6} - \frac{9}{2a+12} = 3$

59. $\frac{9}{4a^2-1} - \frac{6}{2a-1} = \frac{4}{2a+1}$

60. $\frac{8}{6n^2-13n-5} + \frac{6}{10n^2-25n} = 0$

Solve the given equations for the indicated variable.

61. $\frac{2+m}{6} - \frac{n-3}{4} = \frac{p}{8}$ for p

62. $F = \frac{9}{5}C + 32$ for C (Temperature conversion)

63. $\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$ for V_1 (Physics formula)

64. In electricity, the total resistance R_t for two resistors connected in parallel is given by

$$R_t = \frac{R_1R_2}{R_1 + R_2}$$

Solve for R_2 .

[4-8]

Set up an equation to solve the following problems.

65. If Erin can do a job in 3 days and Sarah takes 6 days to do the same job, how long would it take the girls to do the job working together?

66. A train travels 120 miles in the same time that an automobile travels 80 miles. If the train travels 30 miles per hour faster than the automobile, what is the speed of each?

67. A boat travels 14 miles downstream in the same amount of time that it takes to travel 10 miles upstream. If the boat travels at 18 miles per hour in still water, what is the speed of the current?

68. The reciprocal of four times a number is equal to the reciprocal of twice the number added to 7. Find the number.

Chapter 4 cumulative testEvaluate the following expressions when $x = 2$ and $y = -3$.

[1-5] 1. $x^2 - 2xy + y^2$

[1-5] 2. $3(2x - 1) - 2(3x + 2)$

[1-5] 3. $\frac{3x^2}{5y^2}$

[4-4] 4. $\frac{\frac{1}{x} - \frac{1}{y}}{\frac{1}{x} + \frac{1}{y}}$

[1-5] 5. $\frac{2}{3x} - \frac{4}{5y} + \frac{1}{xy}$

Perform the indicated operations and simplify. Assume all denominators are nonzero.

[1-4] 6. $3(2x - 1) + 4(x + 5) - (3x + 6)$

[3-3] 7. $(4ab)(-2a^2b)(3a^3b^3)$

[3-2] 8. $(x + 4)(x^2 - 2x + 5)$

[4-2] 9. $\frac{10xy}{9x^2y} \cdot \frac{27x^3y^2}{15x^2y^2}$

[4-2] 10. $\frac{x^2 + 5x - 24}{x^2 - 2x} \div \frac{x^2 - 9}{x^2 - 4x + 4}$

[4-3] 11. $\frac{2y - 3}{8} - \frac{4y + 1}{6}$

[3-2] 12. $(5x - 3)(2x + 9)$

[3-2] 13. $(4x - 5)^2$

[3-2] 14. $(3y - 5)(3y + 5)$

[4-5] 15. $(2x^3 + 3x^2 - 1) \div (x - 3)$

[4-2] 16. $\frac{6a^2 - a - 2}{a^2 + 4a - 21} \cdot \frac{2a^2 - a - 15}{8a^2 - 6a - 5}$

[4-3] 17. $\frac{4}{2x^2 - x - 3} - \frac{5}{2x^2 + 3x + 1}$

State the domain of the given rational expression.

[4-1] 18. $\frac{4y + 1}{2y - 3}$

[4-1] 19. $\frac{3x}{4x^2 - 1}$

[4-1] 20. $\frac{x - 5}{x^2 - 25}$

Find the solution set of each equation.

[2-1] 21. $4(y + 3) - 3(3y + 1) + (4y + 5) = 0$

[4-7] 22. $\frac{3}{4x} - 2 = \frac{5}{3x}$

[4-7] 23. $\frac{3}{y + 2} + \frac{2y + 1}{y^2 - 4} = \frac{-3}{y - 2}$

Simplify each complex rational expression. Assume all denominators are nonzero.

[4-4] 24. $\frac{\frac{1}{4} + \frac{2}{5}}{\frac{3}{10} - 1}$

[4-4] 25. $\frac{4 - \frac{3}{x + 2}}{5 + \frac{2}{x - 2}}$

Find the solution set of the following inequalities.

[2-5] 26. $5 - 3x \geq 8$

[2-5] 27. $3(4y - 1) < 2(5y + 7)$

[2-5] 28. $\frac{z + 1}{4} - \frac{z - 3}{6} > \frac{1}{3}$

[2-5] 29. $\frac{3}{4}(x + 1) - 2(x - 3) \leq \frac{2}{3}(x - 2)$

Reduce the following rational expressions to lowest terms. Assume all denominators are nonzero.

[4-1] 30. $\frac{36a^2b^3}{-24ab^5}$

[4-1] 31. $\frac{p^2 + p - 12}{p^2 - 9}$

[4-1] 32. $\frac{12 - 6y}{4y - 8}$

[4-7] 33. Find the solution set of the rational equation

[4-8] 34. The sum of twice a number and six times its reciprocal is 13. Find the number.

$$\frac{4}{x} - 3 = 6 + \frac{5}{x}$$

[4-6] 35. Using synthetic division, find $P(-2)$ when $P(x) = x^4 - x^3 + 2x^2 - 5$.

Exercise 3–8

Answers to odd-numbered problems

1. $(m-7)(m+7)$ 3. $(x+5)(x+1)$ 5. $(7a+1)(a+5)$
7. $(2a+3)(a+6)$ 9. $(ab+4)(ab-2)$
11. $(3a+b)(9a^2-3ab+b^2)$ 13. $5(3x+y)(5x^2+a)$
15. $10(x-y)^2$ 17. $4(m-2n)(m+2n)$ 19. $(a-b-2x-y)(a-b+2x+y)$
21. $3(x^2-3y)(x^4+3x^2y+9y^2)$
23. $2xy^2(6x^2-9x+8y^2)$ 25. $(2x+3y)(2x-3y)$
27. $(x+2y)(3a-b)$ 29. $(3a^3-bc)(9a^6+3a^3bc+b^2c^2)$
31. $(5a+3)(a-7)$ 33. $(a-1)(a+1)(a-2)(a+2)$
35. $(2a+3b)(2a-5b)$ 37. $(y-2)(y+2)(y^2+4)$
39. $2(a+2)(2a+1)$ 41. $(x+y-9)(x+y+1)$
43. $(2a-1)(3a+5)$ 45. $4ab(x+3y)(1-2ab)$
47. $(2a-5b)^2$ 49. $5x(4x^2+1)(2x+1)(2x-1)$
51. $3ab(a^2-3b^2)^2$ 53. $(3a-x-5y)(3a+x+5y)$
55. $(3x-13)(x+7)$ 57. $3x^2(xy^3+3z^2)(x^2y^6-3xy^3z^2+9z^4)$
59. $(3-x)(3+x)(a-3)^2$

Review exercises

1. $\{7\}$ 2. $\left\{-\frac{3}{2}\right\}$ 3. $\left\{\frac{4}{5}\right\}$ 4. $\{0\}$ 5. $\left\{\frac{6}{5}\right\}$ 6. $\{2\}$

Chapter 3 review

1. $-15x^5$ 2. $6a^3b^5$ 3. $8a^2b^{12}c^3$ 4. $x^8y^9z^2$ 5. $17a^7$
6. $4b^{10}-27b^8$ 7. $10a^4b-15a^3b^2+20a^2b^3$ 8. $2a^2+ab-b^2$
9. y^2-49 10. $4a^2+4ab+b^2$ 11. $a^3-2a^2b-ab^2+2b^3$
12. $2x^2+16x-17$ 13. $\frac{b^3}{4a^3}$ 14. $\frac{6}{x^9}$ 15. $\frac{9y^6}{x^4}$ 16. $\frac{64x^9}{y^{15}}$
17. $\frac{a^9}{b^3}$ 18. $\frac{a^3c^7}{2b}$ 19. $\frac{4}{y^5}$ 20. $6x^2(2x-3)$
21. $4a^2b(3a^2-b^2+6ab)$ 22. $(x-2y)(a+b)$
23. $5x(2x+1)(x-3z)$ 24. $(2x-y)(3a-b)$
25. $(2y+x)(4a+3b)$ 26. $(2x+y)(2a+3b)$
27. $(a+3b)(x-2y)$ 28. $(2x-y)(3a-b)$
29. $(a+3b)(2x+y)$ 30. $(a+2)(a+12)$
31. $(b-7)(b-2)$ 32. $2a(a-5)(a+1)$
33. $x(x-3)(x+2)$ 34. $(xy+6)(xy+4)$
35. $(ab-10)(ab+2)$ 36. $(x+3y-1)(x+3y+2)$
37. $(x+2y-7)^2$ 38. $(a-2b)(a+b)$
39. $(x+2y)(x+3y)$ 40. $(2a-3)(a-2)$
41. $(4x-5)(2x-1)$ 42. $(2y+1)(3y-4)$
43. $(3a-1)(a+1)$ 44. $(4x-1)(x+3)$
45. $(2x+1)(2x+1)=(2x+1)^2$ 46. $(6a+1)(4a+3)$
47. $(4x-3)(2x-3)$ 48. $(2x+3)(x+6)$
49. $3(2x+1)(x+8)$ 50. $(-2x-1)(x-6)$ or $(2x+1)(-x+6)$ 51. $(x+9)(x-9)$
52. $4(a+3b)(a-3b)$ 53. $3(a+3b)(a-3b)$
54. $(y^2+9)(y+3)(y-3)$ 55. $(a+2b)(x+y)(x-y)$
56. $(x-3z)(2a+3b)(2a-3b)$ 57. $(a+2b)(a^2-2ab+4b^2)$
58. $(3x-y)(9x^2+3xy+y^2)$ 59. $8(2a+b)(4a^2-2ab+b^2)$
60. $x^2(3x+y)(9x^2-3xy+y^2)$ 61. $2(x-2b)(x^2+2bx+4b^2)$
62. $(x^4y^5-z^3)(x^8y^{10}+x^4y^5z^3+z^6)$ 63. $(3x+4)^2$
64. $(a^2-b^3)(a^4+a^2b^3+b^6)$ 65. $3a(a+5)(a-5)$
66. $(a+6)^2(3+x)(3-x)$ 67. $(6a-1)(a+3)$
68. $(x-2y)(4m+3n)$ 69. $5x(2x+y)(4x^2-2xy+y^2)$
70. $ab^3(a-2b)^2$ 71. $(x+4y)(x-2y)$
72. $(3a+2)(5a-2)$

Chapter 3 cumulative test

1. false 2. false 3. true 4. 2 5. undefined 6. 4 7. 0
8. 81 9. 37 10. -34 11. $\{x|x \geq 7\}$ 12. $\{5\}$
13. $\left\{-3, \frac{1}{3}\right\}$ 14. $\left\{x|-\frac{1}{3} < x < \frac{17}{3}\right\}$
15. $\left\{x|x < \frac{-11}{6} \text{ or } x > \frac{1}{6}\right\}$ 16. $\left\{x|-\frac{3}{2} < x < \frac{7}{2}\right\}$
17. $\left\{x|-5 \leq x \leq \frac{-3}{2}\right\}$ 18. let $x =$ the number; $\{x|x > 9\}$
19. \$4,000 20. $6a^5b^5$ 21. $16x^{12}y^{16}$
22. $\frac{3a^3b^5}{2c^6}$ 23. $\frac{y^4}{x^6}$ 24. $\frac{y^9}{27x^9}$ 25. $4x^2-2x-3$ 26. $\frac{a^9c^3}{27b^6}$
27. $7ab-a-5b$ 28. $9a^2-6ab+b^2$ 29. $8a^{18}b^{11}$
30. $12x^6y^3-18x^6y^4+24x^4y^5$ 31. $(x+6)(x-6)$
32. $(x+3y)(x^2-3xy+9y^2)$ 33. $(2a-3)(a-6)$
34. $(y+6)(y+1)$ 35. $(xy+2)(xy-4)$
36. $(7x+1)(x-5)$ 37. $(2a+3b)(x-3y)$
38. $5x(2a-b)(3x+1)$ 39. $(2x-5y)^2$
40. $3(a+2b)(a-2b)$

Chapter 4

Exercise 4–1

Answers to odd-numbered problems

1. domain = $\{x|x \in R, x \neq 4\}$ 3. domain = $\{x|x \in R, x \neq -1\}$
5. domain = $\left\{z|z \in R, z \neq \frac{3}{2}\right\}$ 7. domain = $\left\{x|x \in R, x \neq -\frac{4}{7}\right\}$
9. domain = $\{x|x \in R, x \neq 0, 4\}$ 11. domain = $\{x|x \in R, x \neq -4\}$
13. domain = $\left\{y|y \in R, y \neq -\frac{5}{2}, \frac{5}{2}\right\}$
15. domain = $\left\{x|x \in R, x \neq -\frac{1}{2}, 3\right\}$
17. domain = $\{x|x \in R\}$ 19. $\frac{5}{3x}$ ($x \neq 0$) 21. $\frac{p^2}{q^2}$ ($p \neq 0, q \neq 0$)
23. $-\frac{ab^3}{c^4}$ ($a \neq 0, b \neq 0, c \neq 0$) 25. $\frac{3n^2p^2}{4m^3}$ ($m \neq 0, n \neq 0, p \neq 0$)
27. $\frac{3}{7}$ ($x \neq 2$) 29. $\frac{3}{a-2}$ ($a \neq 0, 2$) 31. $2(x-1)$ ($x \neq 0$)
33. $\frac{4(y+1)}{3}$ ($y \neq 1$) 35. -5 ($x \neq y$) 37. $\frac{a-3}{4}$ ($a \neq -3$)
39. $2y-1$ ($y \neq -\frac{1}{2}$) 41. x^2-2x+4 ($x \neq -2$)
43. $\frac{-3}{2(y^2+xy+x^2)}$ ($x \neq y$) 45. $\frac{y-7}{y+7}$ ($y \neq -7$)
47. $\frac{m-6}{m-3}$ ($m \neq -2, 3$) 49. $\frac{y-8}{y-5}$ ($y \neq -4, 5$)
51. $\frac{a-1}{a+1}$ ($a \neq -1, \frac{1}{2}$) 53. $\frac{2x-3}{4x+1}$ ($x \neq -3, -\frac{1}{4}$)
55. $\frac{3m+2}{2m+1}$ ($x \neq -\frac{3}{4}, -\frac{1}{2}$) 57. $\frac{2(a+2)}{3a+1}$ ($a \neq -\frac{1}{3}, \frac{3}{2}$)
59. $\frac{5(y+1)}{4(y+3)}$ ($y \neq -3, 3$) 61. $\frac{-(a+6)}{3+a}$ ($a \neq -3, \frac{2}{3}$)
63. $\frac{p+4q}{p+2q}$ ($p \neq -3q, -2q$)

Solutions to trial exercise problems

5. Since $2z - 3 = 0$ when $z = \frac{3}{2}$, the domain is $\left\{z \mid z \in \mathbb{R}, z \neq \frac{3}{2}\right\}$.

10. Since $3m^2 + 6m = 3m(m + 2)$ and $3m = 0$ when $m = 0$, $m + 2 = 0$ when $m = -2$, the domain is $\{m \mid m \in \mathbb{R}, m \neq 0, -2\}$.

13. Since $4y^2 - 25 = (2y + 5)(2y - 5)$ and $2y + 5 = 0$

when $y = -\frac{5}{2}$, $2y - 5 = 0$ when $y = \frac{5}{2}$, the domain is

$\left\{y \mid y \in \mathbb{R}, y \neq -\frac{5}{2}, \frac{5}{2}\right\}$. 17. Since $x^2 + 4 \neq 0$ for any value of x , the domain is $\{x \mid x \in \mathbb{R}\}$.

$$30. \frac{-36x}{42x^3 + 24x} = \frac{-36x}{6x(7x^2 + 4)} = \frac{6x \cdot (-6)}{6x(7x^2 + 4)} = \frac{-6}{7x^2 + 4} \quad (x \neq 0)$$

$$35. \frac{5x - 5y}{y - x} = \frac{5(x - y)}{-(x - y)} = \frac{5}{-1} = -5 \quad (x \neq y)$$

$$41. \frac{x^3 + 8}{x + 2} = \frac{(x + 2)(x^2 - 2x + 4)}{x + 2} = x^2 - 2x + 4 \quad (x \neq -2)$$

$$53. \frac{2x^2 + 3x - 9}{4x^2 + 13x + 3} = \frac{(2x - 3)(x + 3)}{(x + 3)(4x + 1)} = \frac{2x - 3}{4x + 1}$$

$$\left(x \neq -3, -\frac{1}{4}\right) \quad 63. \frac{p^2 + 7pq + 12q^2}{p^2 + 5pq + 6q^2} = \frac{(p + 3q)(p + 4q)}{(p + 2q)(p + 3q)} = \frac{p + 4q}{p + 2q} \quad (p \neq -3q, -2q)$$

Review exercises

1. $-\frac{1}{12}$ 2. 1 3. $\frac{13}{8}$ or $1\frac{5}{8}$ 4. $\{2\}$ 5. $\left\{\frac{9}{2}\right\}$

6. $x = \frac{2a - 5}{2}$

Exercise 4-2

Answers to odd-numbered problems

1. $\frac{8}{3x} \quad (x \neq 0)$ 3. $\frac{1}{2x^2y^2} \quad (x, y \neq 0)$ 5. $\frac{a^2}{7bc^3} \quad (a, b, c \neq 0)$

7. $2ab^2 \quad (a, b \neq 0)$ 9. $\frac{4a - 8}{15a - 60} \quad (a \neq 4, -2)$ 11. $\frac{a - 6}{6} \quad (a \neq -3)$

13. $\frac{-4p^2 - 5p - 1}{p - 1} \quad \left(p \neq -1, \frac{1}{4}, 1\right)$ 15. $\frac{x - 5}{x - 3} \quad (x \neq 1, 2, 3, 4)$

17. $\frac{3a^2 - 5a - 2}{3a^2 - 4a + 1} \quad \left(a \neq -2, -\frac{3}{2}, \frac{1}{3}, 1\right)$

19. $\frac{x^2 - 2x + 4}{2x + 7} \quad \left(x \neq -2, -\frac{7}{2}, 2\right)$ 21. $\frac{-1}{x + 2y} \quad (x \neq -2y, -y, y)$

23. $\frac{a^2}{b^3}$ 25. $\frac{1}{r^2 - 5r + 4}$ 27. $\frac{1}{3x + 12}$ 29. $\frac{x^3 + x^2 - x - 1}{4}$

31. $\frac{x^3 - 125}{x + 5}$ 33. $\frac{-x^2 - 3x + 40}{x^2 + 4x - 21}$ 35. $\frac{x + 4}{x - 4}$ 37. $\frac{a - b}{2}$

39. $-(2m^2 + 5mn + 3n^2)$ 41. $\frac{a - 6}{a + 3}$ 43. $\frac{x^2 - 9}{x^2 - 5x + 4}$

45. $-(9y^2 + 6xy + 4x^2)(x^2 - xy + y^2)$ 47. $\frac{b^2 + b - 2}{b^2 - b - 2}$

49. $\frac{z^2 - 2wz + w^2}{z^2 + 2wz + w^2}$

Solutions to trial exercise problems

9. $\frac{4a + 8}{3a - 12} \cdot \frac{a - 2}{5a + 10} = \frac{4(a + 2) \cdot (a - 2)}{3(a - 4) \cdot 5(a + 2)} = \frac{4(a - 2)}{15(a - 4)}$

$= \frac{4a - 8}{15a - 60} \quad (a \neq -2, a \neq 4)$ 17. $\frac{2a^2 - a - 6}{3a^2 - 4a + 1} \cdot \frac{3a^2 + 7a + 2}{2a^2 + 7a + 6}$

$= \frac{(2a + 3)(a - 2)}{(3a - 1)(a - 1)} \cdot \frac{(3a + 1)(a + 2)}{(2a + 3)(a + 2)} = \frac{(a - 2)(3a + 1)}{(3a - 1)(a - 1)}$

$= \frac{3a^2 - 5a - 2}{3a^2 - 4a + 1} \quad \left(a \neq -2, -\frac{3}{2}, \frac{1}{3}, 1\right)$

25. $\frac{r + 4}{r^2 - 1} \div \frac{r^2 - 16}{r + 1} = \frac{(r + 4) \cdot (r + 1)}{(r^2 - 1) \cdot (r^2 - 16)}$

$= \frac{(r + 4)(r + 1)}{(r + 1)(r - 1)(r + 4)(r - 4)} = \frac{1}{(r - 4)(r - 1)}$

$= \frac{1}{r^2 - 5r + 4}$ 33. $\frac{56 - x - x^2}{x^2 - 6x - 7} \div \frac{x^2 + 4x - 21}{x^2 - 4x - 5}$

$= \frac{-(x^2 + x - 56) \cdot (x^2 - 4x - 5)}{(x^2 - 6x - 7) \cdot (x^2 + 4x - 21)}$

$= \frac{-(x - 7)(x + 8) \cdot (x - 5)(x + 1)}{(x - 7)(x + 1) \cdot (x + 7)(x - 3)} = \frac{-(x + 8)(x - 5)}{(x + 7)(x - 3)}$

$= \frac{-(x^2 + 3x - 40)}{x^2 + 4x - 21} = \frac{-x^2 - 3x + 40}{x^2 + 4x - 21}$

39. $\frac{n^2 - m^2}{2m - 3n} \div \frac{m - n}{4m^2 - 9n^2} = \frac{(n^2 - m^2) \cdot (4m^2 - 9n^2)}{(2m - 3n) \cdot (m - n)}$

$= \frac{-(m - n)(m + n) \cdot (2m + 3n)(2m - 3n)}{(2m - 3n) \cdot (m - n)}$

$= -(m + n)(2m + 3n) = -(2m^2 + 5mn + 3n^2)$

41. $\frac{a^2 - 8a + 15}{a^2 + 7a + 6} \cdot \frac{a^2 - 6a - 7}{a^2 - 9} \div \frac{a^2 - 12a + 35}{a^2 - 36}$

$= \frac{(a^2 - 8a + 15) \cdot (a^2 - 6a - 7) \cdot (a^2 - 36)}{(a^2 + 7a + 6) \cdot (a^2 - 9) \cdot (a^2 - 12a + 35)}$

$= \frac{(a - 3)(a - 5) \cdot (a - 7)(a + 1) \cdot (a + 6)(a - 6)}{(a + 6)(a + 1) \cdot (a + 3)(a - 3) \cdot (a - 7)(a - 5)} = \frac{a - 6}{a + 3}$

47. $\frac{ab - a + 3b - 3}{ab + a - 4b - 4} \div \frac{ab - 2a + 3b - 6}{ab + 2a - 4b - 8}$

$= \frac{(ab - a + 3b - 3) \cdot (ab + 2a - 4b - 8)}{(ab + a - 4b - 4) \cdot (ab - 2a + 3b - 6)}$

$= \frac{(a + 3)(b - 1) \cdot (a - 4)(b + 2)}{(a - 4)(b + 1) \cdot (a + 3)(b - 2)} = \frac{(b - 1)(b + 2)}{(b + 1)(b - 2)}$

$= \frac{b^2 + b - 2}{b^2 - b - 2}$

Review exercises

1. $\frac{43}{24}$ or $1\frac{19}{24}$ 2. $\frac{1}{3}$ 3. $\{y \mid y \leq 2\} = (-\infty, 2]$

4. $\{y \mid -3 \leq y < 1\} = [-3, 1)$ 5. $8x^5 - 4x^4 + 24x^3$

6. $4y^2 - 15y - 4$

Exercise 4-3

Answers to odd-numbered problems

1. $70x$ 3. $80k^2$ 5. $288a^3b^2$ 7. $30x^2y^4$ 9. $4a(a - 2)$

11. $6x(x - 7)$ 13. $3(p + 3)^2(p - 4)$ 15. $10(a + 5)(a - 5)$

17. $(q + 7)(q - 7)(q - 2)$ 19. $5(a + b)(a - b)$

21. $(2a + 1)(3a - 1)(a - 7)(a + 7)$ 23. $\frac{12x^2}{21x^3}$ 25. $\frac{15x^3y}{24x^2y^2}$

$$\begin{aligned}
 27. & \frac{4p^2 - 12p}{p - 3} & 29. & \frac{p^2 - 5p + 6}{p^2 - 4} & 31. & \frac{16x^3 + 24x^2 + 36x}{8x^3 - 27} \\
 33. & \frac{2n^2 - 11n + 5}{n^2 + 2n - 35} & 35. & \frac{2y^2 - y - 10}{4y^2 + 7y - 2} & 37. & \frac{15m^2 + 12m}{25m^2 - 16} \\
 39. & \frac{9}{a - 9} & 41. & \frac{13}{q} & 43. & \frac{-3y}{y + 4} & 45. & 5 & 47. & 1 & 49. & \frac{1}{x + 3} \\
 51. & 1 & 53. & \frac{47}{12x} & 55. & \frac{5z - 2}{z^2} & 57. & \frac{18x + 5}{9x^2} & 59. & \frac{6y - 4x}{9x^2y^2} \\
 61. & \frac{14a^2 - 2a}{(3a + 5)(2a - 3)} & 63. & \frac{129y - 122}{10(y - 2)(y + 2)} & 65. & \frac{2x - 4}{x - 9} \\
 67. & \frac{16x - 61}{(x - 6)(x + 1)(x - 1)} & 69. & \frac{5x^2 + 3xy + 6y^2}{(x - 3y)(x + y)^2} \\
 71. & \frac{2a^2 - 16a - 25}{(2a - 1)(4a + 3)(a + 5)} & 73. & \frac{20a^2 - 25a + 1}{5a - 2} \\
 75. & \frac{17p^2 - 20p - 75}{8p(p - 5)(p + 3)} & 77. & \frac{3y^2 - y + 15}{(y + 3)(y - 2)(y + 2)} \\
 79. & \frac{5b^2 + 7b - 30}{5(b + 2)(b - 2)} & 81. & \frac{3x^2 - 15xy + 14y^2}{(x - 6y)^2(x + 2y)} \\
 83. & \frac{15x^2 - 9x - 9}{(4x - 3)(2x - 5)(3x + 1)} \\
 85. & \frac{49m^2 - 20mn + 3n^2}{(8m - n)(m + 2n)(5m - 4n)} & 87. & \frac{qrs + prs + pqs + pqr}{pqrs} \\
 89. & \frac{f_1 + f_2 - d}{f_1 f_2} & 91. & \frac{(n - 1)(R_2 + R_1)}{R_1 R_2}
 \end{aligned}$$

Solutions to trial exercise problems

$$13. \quad p^2 - p - 12 = (p - 4)(p + 3); \quad p^2 + 6p + 9 = (p + 3)^2; \quad 3p - 12 = 3(p - 4). \text{ The LCM is } 3(p + 3)^2(p - 4).$$

$$18. \quad 2y + 10 = 2(y + 5); \quad 25 - y^2 = -(y + 5)(y - 5); \quad y^2 - 10y + 25 = (y - 5)^2. \text{ The LCM is } 2(y + 5)(y - 5)^2.$$

$$21. \quad 2a^2 - 13a - 7 = (2a + 1)(a - 7); \quad 6a^2 + a - 1 = (3a - 1)(2a + 1); \quad a^2 - 49 = (a + 7)(a - 7). \text{ The LCM is } (2a + 1)(3a - 1)(a - 7)(a + 7).$$

$$31. \quad \text{Since } 8x^3 - 27 = (2x - 3)(4x^2 + 6x + 9), \text{ then } \frac{4x}{2x - 3} = \frac{4x \cdot (4x^2 + 6x + 9)}{(2x - 3) \cdot (4x^2 + 6x + 9)} = \frac{16x^3 + 24x^2 + 36x}{8x^3 - 27}. \quad 37. \quad \text{Since}$$

$$25m^2 - 16 = (5m + 4)(5m - 4), \text{ then } -\frac{3m}{4 - 5m} = -\frac{3m}{-(5m - 4)} = \frac{3m}{5m - 4} = \frac{3m \cdot (5m + 4)}{(5m - 4)(5m + 4)} = \frac{15m^2 + 12m}{25m^2 - 16}.$$

$$53. \quad \frac{8}{3x} + \frac{5}{4x} = \frac{8(4)}{3x(4)} + \frac{5(3)}{4x(3)} = \frac{32}{12x} + \frac{15}{12x} = \frac{32 + 15}{12x} = \frac{47}{12x}$$

$$65. \quad \frac{x + 2}{x - 9} - \frac{x - 6}{9 - x} = \frac{x + 2}{x - 9} - \frac{x - 6}{-(x - 9)} = \frac{x + 2}{x - 9} + \frac{x - 6}{x - 9} = \frac{(x + 2) + (x - 6)}{x - 9} = \frac{x + 2 + x - 6}{x - 9} = \frac{2x - 4}{x - 9}$$

$$70. \quad \frac{8q}{4q^2 - 9p^2} + \frac{5q}{4q^2 - 12pq + 9p^2} = \frac{8q}{(2q + 3p)(2q - 3p)} + \frac{5q}{(2q - 3p)^2} = \frac{8q(2q - 3p)}{(2q + 3p)(2q - 3p)^2} + \frac{5q(2q + 3p)}{(2q - 3p)^2(2q + 3p)} = \frac{(16q^2 - 24pq) + (10q^2 + 15pq)}{(2q + 3p)(2q - 3p)^2} = \frac{26q^2 - 9pq}{(2q + 3p)(2q - 3p)^2}$$

$$\begin{aligned}
 73. & (4a - 3) - \frac{2a + 5}{5a - 2} = \frac{4a - 3}{1} - \frac{2a + 5}{5a - 2} \\
 & = \frac{(4a - 3) \cdot (5a - 2) - 1 \cdot (2a + 5)}{1 \cdot (5a - 2)} = \frac{20a^2 - 23a + 6 - 2a - 5}{5a - 2} = \frac{20a^2 - 25a + 1}{5a - 2} \\
 82. & \frac{a - b}{a^2 - 3ab - 4b^2} + \frac{2b - 5a}{a^2 - 16b^2} = \frac{a - b}{(a - 4b)(a + b)} + \frac{2b - 5a}{(a - b) \cdot (a + 4b)} \\
 & = \frac{(a - b)(a + 4b) + (2b - 5a)(a + b)}{(a - 4b)(a + b)(a + 4b)} \\
 & = \frac{(a^2 + 3ab - 4b^2) + (2b^2 - 3ab - 5a^2)}{(a - 4b)(a - b)(a + b)} \\
 & = \frac{a^2 + 3ab - 4b^2 + 2b^2 - 3ab - 5a^2}{(a - 4b)(a - b)(a + b)} = \frac{-4a^2 - 2b^2}{(a - 4b)(a - b)(a + b)}
 \end{aligned}$$

Review exercises

$$\begin{aligned}
 1. & (2x^2 + 3)(4x - 1) & 2. & (2x + 1)(2x - 7) \\
 3. & (3y + 7)(3y - 7) & 4. & 12 & 5. & 24k
 \end{aligned}$$

Exercise 4-4

Answers to odd-numbered problems

$$\begin{aligned}
 1. & \frac{21}{20} & 3. & \frac{11}{3} & 5. & \frac{12}{11} & 7. & \frac{7}{8m} & 9. & \frac{-4y}{3y + 3} & 11. & \frac{x - 3}{x + 7} \\
 13. & \frac{ab}{b - a} & 15. & \frac{x}{x - 2y} & 17. & \frac{20}{27} & 19. & \frac{5}{7y} & 21. & \frac{35 - 7a}{4a + 36} \\
 23. & \frac{8x - 4y}{3} & 25. & \frac{4x^2 + 5x - 9}{5x^2 + 16x + 3} & 27. & \frac{xy^2 + x^2}{y^2 - xy^2} \\
 29. & \frac{mn(m + n)}{n - m} & 31. & \frac{5q + 4p^2}{p^2q(p - q)} & 33. & \frac{a^2 + 9a + 23}{a^2 + 7a + 7} \\
 35. & \frac{4y^3 + 17y^2 - 18y - 15}{4y^3 + 25y^2 + 31y - 39} & 37. & \frac{t^2 - 3t - 4}{t^2 + 3t - 18} & 39. & \frac{-3}{2x + 14} \\
 41. & \frac{5b^2 + 24b - 5}{4b - 22} & 43. & \frac{1}{2} & 45. & \frac{3c + 4a - 2b}{5} & 47. & \frac{y + x}{y - x} \\
 49. & \frac{y}{y + x} & 51. & \frac{q^2 + p^2}{p^2} & 53. & \frac{pq}{q - p} & 55. & \frac{V_1 C - it}{RC} \\
 57. & CP = \frac{T_1}{T_2 - T_1} & 59. & r = \frac{2dr_1r_2}{dr_2 + dr_1} = \frac{2r_1r_2}{r_2 + r_1}
 \end{aligned}$$

Solutions to trial exercise problems

$$\begin{aligned}
 25. & \frac{4 - \frac{3}{x + 3}}{5 + \frac{6}{x - 1}} = \frac{4(x + 3)(x - 1) - \frac{3}{x + 3} \cdot (x + 3)(x - 1)}{5(x + 3)(x - 1) + \frac{6}{x - 1} \cdot (x + 3)(x - 1)} \\
 & = \frac{4(x^2 + 2x - 3) - 3(x - 1)}{5(x^2 + 2x - 3) + 6(x + 3)} \\
 & = \frac{4x^2 + 8x - 12 - 3x + 3}{5x^2 + 10x - 15 + 6x + 18} = \frac{4x^2 + 5x - 9}{5x^2 + 16x + 3} \\
 28. & \frac{\frac{1}{a} - \frac{1}{b}}{\frac{1}{a^2} + \frac{1}{b^2}} = \frac{\frac{1}{a} \cdot \frac{a^2b^2}{a^2b^2} - \frac{1}{b} \cdot \frac{a^2b^2}{a^2b^2}}{\frac{1}{a^2} \cdot \frac{a^2b^2}{a^2b^2} + \frac{1}{b^2} \cdot \frac{a^2b^2}{a^2b^2}} = \frac{ab^2 - a^2b}{b^2 + a^2}
 \end{aligned}$$

$$33. \frac{(a+5) + \frac{3}{a+4}}{(a+3) - \frac{5}{a+4}} = \frac{(a+5)(a+4) + \frac{3}{a+4}(a+4)}{(a+3)(a+4) - \frac{5}{a+4}(a+4)}$$

$$= \frac{a^2 + 9a + 20 + 3}{a^2 + 7a + 12 - 5} = \frac{a^2 + 9a + 23}{a^2 + 7a + 7}$$

$$39. \frac{\frac{x^2 - x - 6}{2} - \frac{4}{x+2}}{\frac{x-3}{x+2} - \frac{4}{x-3}} = \frac{\frac{(x-3)(x+2)}{2} - \frac{4}{x+2}}{\frac{(x-3)(x+2)}{x+2} - \frac{4}{x-3}}$$

$$= \frac{\frac{(x-3)(x+2) - 8}{2}}{\frac{(x-3)(x+2) - 8}{x+2}}$$

$$= \frac{(x-3)(x+2) - 8}{2} \cdot \frac{x+2}{(x-3)(x+2) - 8}$$

$$= \frac{2(x-3) - 4(x+2)}{2} = \frac{2x - 6 - 4x - 8}{2} = \frac{-2x - 14}{2} = -x - 7$$

Review exercises

1. $\left\{-\frac{14}{3}\right\}$ 2. $\{-2, 8\}$ 3. $\{2\}$ 4. $\{x | -2 < x < 6\} = (-2, 6)$
 5. $\{x | x \leq -4 \text{ or } x \geq -1\} = (-\infty, -4] \cup [-1, \infty)$
 6. $2a^3 + 8a^2 - a - 4$

Exercise 4–5

Answers to odd-numbered problems

1. $5x^2 - 3x + 2$ 3. $-x^3 + 2x^2 - 3$ 5. $c - 1$
 7. $10xy^2 + 7 - 6y^2$ 9. $3a^4b - 5a^4b^4 + 7a^3b$ 11. $2a + 3c$
 13. $a - 4$ 15. $y + 2 + \frac{1}{y+5}$ 17. $x - 4 - \frac{2}{x-3}$
 19. $a^2 - 2a + 4$ 21. $2y^2 - 3y + 1 + \frac{2}{y+1}$ 23. $x^2 + 3x - 4$
 25. $2a^2 + 3a - 4 + \frac{5}{2a+1}$ 27. $3y^2 - y + 4 + \frac{2}{3y+1}$
 29. $a^2 - 3a + 9$ 31. $3x^3 - 5x^2 + 5x - 4 + \frac{3}{x+1}$
 33. $3a - 1$ 35. $y^2 + y - 1$ 37. $x^2 + 3x + 2 - \frac{4}{x^2 + x - 4}$
 39. $3a^2 - 2a - 4$ 41. $y^2 + 3 + \frac{2}{2y^2 - 3y + 2}$ 43. 2, it is the same. 45. $2x - 3$ 47. $3x - 2$

Solutions to trial exercise problems

$$10. \frac{x(y-2) - z(y-2)}{y-2} = \frac{x(y-2)}{y-2} - \frac{z(y-2)}{y-2} = x - z$$

$$16. (2x - 5 + x^2) \div (x + 4) = (x^2 + 2x - 5) \div (x + 4)$$

$$x + 4 \overline{) x^2 + 2x - 5}$$

$$\underline{x^2 + 4x}$$

$$-2x - 5$$

$$\underline{-2x - 8}$$

$$3$$

Answer: $x - 2 + \frac{3}{x+4}$

$$27. \frac{9y^3 + 11y + 6}{3y + 1} \quad 3y + 1 \overline{) 9y^3 + 0y^2 + 11y + 6}$$

$$\underline{9y^3 + 3y^2}$$

$$-3y^2 + 11y$$

$$\underline{-3y^2 - y}$$

$$12y + 6$$

$$\underline{12y + 4}$$

$$2$$

Answer: $3y^2 - y + 4 + \frac{2}{3y+1}$

$$32. \frac{2x^3 + 5x^2 + 5x + 3}{x^2 + x + 1} \quad x^2 + x + 1 \overline{) 2x^3 + 5x^2 + 5x + 3}$$

$$\underline{2x^3 + 2x^2 + 2x}$$

$$3x^2 + 3x + 3$$

$$\underline{3x^2 + 3x + 3}$$

$$0$$

Answer: $2x + 3$

$$38. \frac{2x^4 - x^3 + 5x^2 - x + 3}{x^2 + 1} \quad x^2 + 0x + 1 \overline{) 2x^4 - x^3 + 5x^2 - x + 3}$$

$$\underline{2x^4 + 0x^3 + 2x^2}$$

$$-x^3 + 3x^2 - x$$

$$\underline{-x^3 - 0x^2 - x}$$

$$3x^2 + 0x + 3$$

$$\underline{3x^2 + 0x + 3}$$

$$0$$

Answer: $2x^2 - x + 3$

Review exercises

1. x 2. x^3 3. $\frac{8}{a^{12}}$ 4. $\frac{3}{10}$ 5. $\frac{-x^2 + 4x + 2}{(x+3)(x-3)}$ 6. $\frac{2x-1}{2x}$
 7. $\left\{\frac{19}{3}\right\}$

Exercise 4–6

Answers to odd-numbered problems

1. $a + 2$ 3. $a + 5$ 5. $x + 7 + \frac{23}{x-2}$ 7. $3a - 1$
 9. $a^2 + 3a + 2 - \frac{2}{a-1}$ 11. $x^2 + x + 1$
 13. $y^2 + 5y + 10 + \frac{16}{y-2}$ 15. $2a^2 + a - 3 + \frac{2}{a-5}$
 17. $x^3 - 2x^2 - x + 3 - \frac{1}{x-3}$ 19. $3y^3 + 2y^2 - 4y - 1 + \frac{3}{y-3}$
 21. $P(1) = -1$ 23. $P(-3) = 319$ 25. $P(0) = 1$
 27. $P(2) = 145$ 29. $P\left(\frac{1}{2}\right) = -\frac{3}{2}$ 31. $P(-3) = 4$;
 $x + 3$ is not a factor 33. $P(2) = 0$; $x - 2$ is a factor
 35. $P(-1) = 25$; $x + 1$ is not a factor 37. $P(-4) = 0$; $x + 4$ is
 a factor 39. $P(-1) = -6$; $x + 1$ is not a factor
 41. $\left\{-8, \frac{2}{3}, 1\right\}$ 43. $\left\{-6, \frac{1}{2}, 5\right\}$ 45. $\{-1, 1\}$
 47. $P(x) = x^3 - 5x^2 + 2x + 8$ 49. $P(x) = x^3 - 3x^2 - 4x$

51. $P(x) = x^4 - 65x^2 + 64$ 53. $P(x) = x^4 - 8x^3 - 13x^2 + 200x - 300$ 55. $P(x) = x^4 - 6x^3 + x^2 + 24x + 16$
 57. $P(x) = x^4 + 7x^3 + 10x^2$ 59. 2 of multiplicity 2; -2 of multiplicity 2; 3 of multiplicity 3; -4 of multiplicity 2 61. 0 of multiplicity 2; -4 of multiplicity 4; 3 of multiplicity 2; 6 63. 25

65. Since the remainder is 0, $\frac{3}{2}$ is a solution.

Solutions to trial exercise problems

$$8. \frac{5 - 3x + 2x^2}{x + 2} = \frac{2x^2 - 3x + 5}{x + 2} = 2x - 7 + \frac{19}{x + 2}$$

$$\begin{array}{r} -2 \overline{2 \quad -3 \quad 5} \\ \downarrow \quad \quad \quad \\ 2 \quad -7 \quad 19 \\ \uparrow \\ x \end{array}$$

$$11. \frac{x^3 - 1}{x - 1} = \frac{x^3 + 0x^2 + 0x - 1}{x - 1} = x^2 + x + 1$$

$$\begin{array}{r} 1 \overline{1 \quad 0 \quad 0 \quad -1} \\ \downarrow \quad \quad \quad \\ 1 \quad 1 \quad 1 \quad 0 \\ \uparrow \quad \uparrow \\ x^2 \quad x \end{array}$$

$$18. \frac{2a^4 + 6a^3 + a^2 + 2a - 5}{a + 3} = 2a^3 + a - 1 - \frac{2}{a + 3}$$

$$\begin{array}{r} -3 \overline{2 \quad 6 \quad 1 \quad 2 \quad -5} \\ \downarrow \quad \quad \quad \\ 2 \quad 0 \quad 1 \quad -1 \quad -2 \\ \uparrow \quad \uparrow \quad \uparrow \\ a^3 \quad a^2 \quad a \end{array}$$

$$23. -3 \overline{3 \quad -2 \quad 1 \quad -4 \quad 1} \\ \downarrow \quad \quad \quad \\ 3 \quad -11 \quad 34 \quad -106 \quad 319$$

$$P(-3) = 319$$

$$28. -2 \overline{8 \quad 0 \quad 0 \quad 0 \quad 3 \quad -2 \quad -5} \\ \downarrow \quad \quad \quad \\ 8 \quad -16 \quad 32 \quad -64 \quad 128 \quad -262 \quad 528$$

$$P(-2) = 523$$

$$35. -1 \overline{1 \quad -9 \quad 18 \quad 0 \quad -3} \\ \downarrow \quad \quad \quad \\ 1 \quad -10 \quad 28 \quad -28 \quad 25 \leftarrow x + 1 \text{ is not a factor}$$

$$40. -2 \overline{1 \quad 7 \quad 4 \quad -12} \\ \downarrow \quad \quad \quad \\ 1 \quad 5 \quad -6 \quad 0$$

$$\text{set } x^2 + 5x - 6 = 0$$

$$(x + 6)(x - 1) = 0$$

$$x + 6 = 0 \quad \text{or} \quad x - 1 = 0$$

$$x = -6 \quad \quad \quad x = 1$$

The solution set is $\{-6, -2, 1\}$.

$$45. x^2 - 2x + 1 \overline{x^4 + 0x^3 - 2x^2 + 0x + 1} \\ \quad \quad \quad \underline{x^4 - 2x^3 + x^2} \\ \quad \quad \quad \quad 2x^3 - 3x^2 + 0x \\ \quad \quad \quad \quad \underline{2x^3 - 4x^2 + 2x} \\ \quad \quad \quad \quad \quad x^2 - 2x + 1 \\ \quad \quad \quad \quad \quad \underline{x^2 - 2x + 1} \\ \quad \quad \quad \quad \quad \quad 0$$

$$x^2 + 2x + 1 = 0$$

$$(x + 1)^2 = 0$$

$$x + 1 = 0$$

$$x = -1$$

The solution set is $\{-1, 1\}$.

47. Since 2, -1 and 4 are zeros, then

$$P(x) = (x - 2)(x + 1)(x - 4) = (x^2 - x - 2)(x - 4) = x^3 - 5x^2 + 2x + 8$$

50. Since -2, 2, 3 and -3 are zeros, then

$$P(x) = (x + 2)(x - 2)(x - 3)(x + 3) = (x^2 - 4)(x^2 - 9) = x^4 - 13x^2 + 36$$

55. Since -1 of multiplicity 2 and 4 of multiplicity 2 are zeros, then

$$P(x) = (x + 1)^2(x - 4)^2 = (x^2 + 2x + 1)(x^2 - 8x + 16) = x^4 - 6x^3 + x^2 + 24x + 16$$

Review exercises

$$1. \emptyset \quad 2. \{24\} \quad 3. \{4\} \quad 4. 9 \text{ and } 36 \quad 5. R = \frac{3V + \pi h^3}{3\pi h^2}$$

Exercise 4-7

Answers to odd-numbered problems

1. $\{-21\}$ 3. $\left\{\frac{-15}{11}\right\}$ 5. $\left\{\frac{10}{3}\right\}$ 7. $\left\{\frac{67}{7}\right\}$ 9. $\{12\}$
 11. $\left\{\frac{-13}{6}\right\}$ 13. \emptyset 15. $\left\{\frac{-3}{5}\right\}$ 17. $\left\{\frac{-13}{5}\right\}$
 19. \emptyset ; 2 is extraneous 21. $\left\{\frac{13}{3}\right\}$ 23. $\left\{\frac{-16}{5}\right\}$ 25. $\left\{\frac{13}{2}\right\}$
 27. \emptyset ; 3 is extraneous 29. $\{-15\}$ 31. $\left\{\frac{-17}{3}\right\}$ 33. $\{26\}$
 35. $\left\{\frac{-31}{5}\right\}$ 37. $b = \frac{3a}{4 - 7a}$ 39. $p = \frac{3q - 3}{3a + 4}$
 41. $y = \frac{5x + 19}{3}$ 43. $r = \frac{S - a}{S}$ 45. $b_1 = \frac{2A - b_2h}{h}$
 47. $t = \frac{L_1 - L_0}{kL_0}$; $L_0 = \frac{L_1}{kt + 1}$ 49. $t = \frac{I}{Pr}$
 51. $R_3 = \frac{RR_1R_2}{R_1R_2 - RR_2 - RR_1}$ 53. $A = \frac{aF}{f}$

$$55. T_2 = \frac{R_2 M + R_2 T_1 - R_1 M}{R_1}; R_1 = \frac{R_2(M + T_1)}{M + T_2} \quad 57. \text{ a. } \frac{1}{10}$$

$$\text{b. } \frac{1}{2} \quad \text{c. } \frac{t}{10} \quad 59. \text{ a. } \frac{200}{5} = 40 \text{ mph} \quad \text{b. } \frac{200}{4} = 50 \text{ mph}$$

$$\text{c. } \frac{200}{t} \text{ mph} \quad 61. \text{ a. } \frac{1}{n}, \frac{3}{n} \quad \text{b. } \frac{15}{n} \quad 63. \frac{5+n}{6-n}$$

Solutions to trial exercise problems

$$5. \frac{4}{a} - \frac{6}{3a} = \frac{3}{5}$$

Multiply by the LCM, $15a$.

$$15a \cdot \frac{4}{a} - 15a \cdot \frac{6}{3a} = 15a \cdot \frac{3}{5}$$

$$15 \cdot 4 - 5 \cdot 6 = 3a \cdot 3$$

$$60 - 30 = 9a$$

$$30 = 9a$$

$$\frac{10}{3} = a$$

The solution set is $\left\{\frac{10}{3}\right\}$.

$$16. 1 + \frac{5}{3m-9} = \frac{10}{m-3}$$

Multiply by the LCM, $3(m-3)$.

$$3(m-3) \cdot 1 + 3(m-3) \cdot \frac{5}{3(m-3)} = 3(m-3) \cdot \frac{10}{m-3}$$

$$3m - 9 + 5 = 3 \cdot 10$$

$$3m - 4 = 30$$

$$3m = 34$$

$$m = \frac{34}{3}$$

The solution set is $\left\{\frac{34}{3}\right\}$.

$$21. 4 - \frac{2x}{5-x} = \frac{6}{x-5}$$

Since $5-x = -(x-5)$, then we have

$$4 - \frac{2x}{-(x-5)} = \frac{6}{x-5}$$

$$4 + \frac{2x}{x-5} = \frac{6}{x-5} \quad \text{Multiply by } x-5.$$

$$4(x-5) + (x-5) \cdot \frac{2x}{x-5} = (x-5) \cdot \frac{6}{x-5}$$

$$4x - 20 + 2x = 6$$

$$6x - 20 = 6$$

$$6x = 26$$

$$x = \frac{26}{6} = \frac{13}{3}$$

The solution set is $\left\{\frac{13}{3}\right\}$.

$$29. \frac{6}{q^2+q-6} = \frac{5}{q^2+3q-10}$$

Since $q^2+q-6 = (q+3)(q-2)$ and $q^2+3q-10 = (q+5)(q-2)$, the LCM is $(q+3)(q-2)(q+5)$.

Multiply each member by the LCM.

$$(q+3)(q-2)(q+5) \cdot \frac{6}{(q+3)(q-2)} = (q+3)(q-2)(q+5) \cdot \frac{5}{(q+5)(q-2)}$$

$$(q+5) \cdot 6 = (q+3) \cdot 5$$

$$6q + 30 = 5q + 15$$

$$q = -15$$

The solution set is $\{-15\}$.

$$41. \frac{y-3}{x+2} = \frac{5}{3}$$

Multiply by the LCM, $3(x+2)$.

$$3(x+2) \cdot \frac{y-3}{x+2} = 3(x+2) \cdot \frac{5}{3}$$

$$3(y-3) = (x+2) \cdot 5$$

$$3y - 9 = 5x + 10$$

$$3y = 5x + 19$$

$$y = \frac{5x+19}{3}$$

$$48. R_x = R_m \left(\frac{E_1}{E_2} - 1 \right)$$

$$\text{Now } R_x = R_m \cdot \frac{E_1}{E_2} - R_m$$

$$R_x + R_m = \frac{R_m E_1}{E_2}$$

Multiply each member by E_2 .

$$E_2(R_x + R_m) = R_m E_1$$

Divide each member by R_m .

$$\frac{E_2(R_x + R_m)}{R_m} = E_1$$

57. If Ann can paint the house in 10 hours, she can paint

$$\text{a. } 1 \div 10 = \frac{1}{10}, \quad \text{b. } 5 \div 10 = \frac{5}{10} = \frac{1}{2}, \quad \text{c. } t \div 10 = \frac{t}{10} \text{ of}$$

the house in 1 hour, 5 hours, and t hours. 61. Let $n = 1$ st number,then $\frac{1}{3}n = \frac{n}{3} = 2$ nd number. a. The reciprocals are $\frac{1}{n}$ and $\frac{3}{n}$.

$$\text{b. Five times the 2nd reciprocal } 5 \cdot \frac{3}{n} = \frac{15}{n}.$$

Review exercises

$$1. (2p+5q)(2p-5q) \quad 2. (x-12)^2 \quad 3. (2y+1)(y-8)$$

$$4. \$12,000 \text{ at } 6\%; \$8,000 \text{ at } 8\% \quad 5. 8 \quad 6. P(-3) = 12$$

Exercise 4-8

Answers to odd-numbered problems

$$1. 1 \text{ hour } 12 \text{ minutes} \quad 3. 40 \text{ minutes} \quad 5. 45 \text{ hours}$$

$$7. 13\frac{11}{13} \text{ hours} \approx 13.8 \text{ hours} \quad 9. 18 \text{ hours for pump } B;$$

$$36 \text{ hours for pump } A \quad 11. 7 \text{ hours} \quad 13. \frac{48}{17} \text{ mph} \quad 15. 30 \text{ km/hr}$$

$$17. 1,600 \text{ miles} \quad 19. 1\frac{2}{3} \text{ miles} \quad 21. \frac{2}{3} \text{ and } 2 \quad 23. 13$$

$$25. 3 \quad 27. 3 \text{ mph} \quad 29. 6 \text{ hours} \quad 31. 30 \text{ mph}$$

$$33. \text{ Sarah, } 4 \text{ mph; Erin, } 6 \text{ mph}$$

Solutions to trial exercise problems

4. Let x = minutes taken to mix the 12 loaves working together.

$$\frac{1}{36} + \frac{1}{40} + \frac{1}{30} = \frac{1}{x} \quad \text{Multiply by the LCM, } 360x.$$

$$\begin{aligned} 360x \cdot \frac{1}{36} + 360x \cdot \frac{1}{40} + 360x \cdot \frac{1}{30} &= 360x \cdot \frac{1}{x} \\ 10x + 9x + 12x &= 360 \\ 31x &= 360 \\ x &= \frac{360}{31} = 11\frac{19}{31} \end{aligned}$$

Working together, they could mix the 12 loaves in $11\frac{19}{31}$ minutes
 ≈ 11.6 minutes.

7. Let x = number of hours necessary to fill the basin with all pipes open.

$$\frac{1}{10} + \frac{1}{12} - \frac{1}{9} = \frac{1}{x} \quad \text{Multiply by the LCM, } 180x.$$

$$\begin{aligned} 180x \cdot \frac{1}{10} + 180x \cdot \frac{1}{12} - 180x \cdot \frac{1}{9} &= 180x \cdot \frac{1}{x} \\ 18x + 15x - 20x &= 180 \\ 13x &= 180 \\ x &= \frac{180}{13} = 13\frac{11}{13} \end{aligned}$$

It would take $13\frac{11}{13}$ hours ≈ 13.8 hours to fill the basin with all three pipes open.

10. Let x = time for slower microprocessor to do the job. Then

$$\frac{3}{5}x = \text{time for faster microprocessor to do the job.}$$

$$\frac{1}{x} + \frac{1}{\frac{3}{5}x} = \frac{1}{2}$$

$$\frac{1}{x} + \frac{5}{3x} = \frac{1}{2}$$

$$6x \cdot \frac{1}{x} + 6x \cdot \frac{5}{3x} = 6x \cdot \frac{1}{2}$$

$$6 + 2 \cdot 5 = 3x$$

$$16 = 3x$$

$$x = \frac{16}{3}$$

$$\frac{3}{5}x = \frac{16}{5}$$

It would take the microprocessors $\frac{16}{3}$ or $5\frac{1}{3}$ milliseconds and

$\frac{16}{5}$ or $3\frac{1}{5}$ milliseconds, respectively, to process the set of inputs individually.

14. Let r = speed of the wind. Then $300 + r$ = speed of the plane with the wind and $300 - r$ = speed of the plane against the wind.

$$\text{Now } t_w (\text{time with the wind}) = \frac{950}{300 + r}$$

$$t_a (\text{time against the wind}) = \frac{650}{300 - r}$$

$$\text{Then } \frac{950}{300 + r} = \frac{650}{300 - r}.$$

Multiply each member by $(300 + r)(300 - r)$.

$$(300 + r)(300 - r) \cdot \frac{950}{300 + r} = (300 + r)(300 - r) \cdot \frac{650}{300 - r}$$

$$(300 - r) \cdot 950 = (300 + r) \cdot 650$$

$$285,000 - 950r = 195,000 + 650r$$

$$90,000 = 1,600r$$

$$r = \frac{90,000}{1,600}$$

$$r = \frac{900}{16} = 56\frac{1}{4}$$

The speed of the wind is $56\frac{1}{4}$ mph.

21. Let n = one number, then $3n$ = the other number. The

reciprocals are $\frac{1}{n}$ and $\frac{1}{3n}$.

$$\begin{array}{ccc} \text{sum of the reciprocals} & \text{is} & 2 \\ \downarrow & & \downarrow \\ \frac{1}{n} + \frac{1}{3n} & = & 2 \end{array}$$

$$3n \cdot \frac{1}{n} + 3n \cdot \frac{1}{3n} = 3n \cdot 2$$

$$3 + 1 = 6n$$

$$4 = 6n$$

$$n = \frac{4}{6} = \frac{2}{3}$$

$$3n = 3 \cdot \frac{2}{3} = 2$$

The numbers are $\frac{2}{3}$ and 2.

27. Let x = speed of current.

	d	r	t
downstream	10	$12 + x$	$\frac{10}{12 + x}$
upstream	6	$12 - x$	$\frac{6}{12 - x}$

Since times are the same,

$$\frac{10}{12 + x} = \frac{6}{12 - x}$$

$$10(12 - x) = 6(12 + x)$$

$$120 - 10x = 72 + 6x$$

$$16x = 48$$

$$x = 3$$

The current has a speed of 3 mph.

30. Inlet pipe fills $\frac{1}{45}$ of the tank in 1 minute

Outlet pipe empties $\frac{1}{30}$ of the tank in 1 minute.

Let x = number of minutes to empty the tank

$$\frac{1}{30} - \frac{1}{45} = \frac{1}{x}$$

$$90x \cdot \frac{1}{30} - 90x \cdot \frac{1}{45} = 90x \cdot \frac{1}{x}$$

$$3x - 2x = 90$$

$$x = 90$$

It would take 90 minutes to empty the tank.

Review exercises

1. $8x^3 - 12x^2 + 4x$ 2. $3x^2 - 16x + 5$ 3. $25z^2 - 40z + 16$

4. $4y^2 - 9$ 5. $\frac{1}{y}$ 6. x^4y^3 7. -7 or 7 8. -5

Chapter 4 review

1. $\{x|x \in R, x \neq -7\}$ 2. $\{x|x \in R, x \neq \frac{4}{3}\}$
3. $\{x|x \in R, x \neq 5\}$ 4. $\{a|a \in R, a \neq -\frac{4}{3}, \frac{4}{3}\}$
5. $\{z|z \in R, z \neq -\frac{5}{2}, \frac{1}{3}\}$ 6. $\{y|y \in R, y \neq \frac{2}{3}\}$ 7. $\frac{ab^3}{c^2}$
8. $\frac{-2n^2}{7mp^4}$ 9. $\frac{5}{6}$ 10. $\frac{3}{a-2}$ 11. $\frac{-5}{2y+x}$
12. $\frac{y^2+4y+16}{y+4}$ 13. $\frac{a-12}{a+1}$ 14. $\frac{4x+3}{5x-1}$
15. $\frac{-(2y+3)}{2(3y+2)}$ 16. $\frac{6y}{x} (x \neq 0, y \neq 0)$
17. $12ay (a \neq 0, y \neq 0)$ 18. $\frac{(4p+3)(p-4)}{3} (p \neq -4, \frac{3}{4})$
19. $\frac{z-3}{2(z+1)(z-1)} (z \neq -1, 1, 3)$
20. $\frac{(m^2+2m+4)(m+6)}{m(m+5)} (m \neq -5, 0, 2, 3)$
21. $1 (a \neq -7, -3, -\frac{2}{5}, \frac{1}{2})$
22. $\frac{(x+7)(2x-1)}{(4x^2-2x+1)(x+2)} (x \neq -2, -\frac{1}{2}, \frac{1}{2}, 7)$
23. $\frac{x^2}{(4x+5)^2} (x \neq -\frac{5}{4}, \frac{5}{4})$ 24. $\frac{y+3}{y+2} (y \neq -3, -2, \frac{4}{7})$
25. $\frac{m-n}{m+n} (m \neq -n, \frac{n}{2}; q \neq -p, p)$ 26. $180x^3y^3$
27. $6x^2(x+2)(x+4)(x-4)$ 28. $3a(a+5)(a-2)$
29. $p(p-5)(p+5)^2$ 30. $\frac{41x}{12y}$ 31. $\frac{-2n^2-28n-25}{(n+4)(n-1)}$
32. $\frac{10p^2+29p+81}{p(p+9)(p+2)(p-2)}$ 33. $\frac{6b^2+16b-23}{3b-2}$
34. $\frac{2y^2+18y+3}{(y+7)(y-7)}$ 35. $\frac{-(4x^2+15x+4)}{(x-6)(x+6)(x+4)}$
36. $\frac{37}{2(a-2)}$ 37. $\frac{-5x^2+63x-102}{8(x-7)(x+4)}$ 38. $\frac{4a+b}{(a-2b)(a+2b)}$
39. $\frac{1}{R_t} = \frac{I_1E_2E_3 + I_2E_1E_3 + I_3E_1E_2}{E_1E_2E_3}$ 40. $\frac{2}{a}$ 41. $\frac{5x}{4x-12}$

42. $\frac{3x+6}{x-5}$ 43. $\frac{7b-38}{8b-34}$ 44. $\frac{x^2y-xy^2}{2y+3x}$
45. $\frac{p^2-4p+5}{p^2-4}$ 46. $5a^6+3a^2+2$ 47. $6a^2b^2c^2-3c^3+1$
48. $3x^2-3x+4$ 49. x^3+x^2-x+1 50. $P(-2) = -35$
51. $P(1) = 11$ 52. $P(-1) = -12$ 53. -3 is a solution
54. -1 is not a solution 55. 2 is not a solution 56. $\left\{\frac{55}{216}\right\}$
57. $\{66\}$ 58. $\left\{-\frac{37}{6}\right\}$ 59. $\left\{\frac{7}{20}\right\}$ 60. $\left\{-\frac{3}{29}\right\}$
61. $p = \frac{4m-6n+26}{3}$ 62. $C = \frac{5}{9}(F-32)$
63. $V_1 = \frac{P_2V_2T_1}{P_1T_2}$ 64. $R_2 = \frac{R_1R_1}{R_1-R_1}$ 65. 2 days
66. 60 mph, automobile; 90 mph, train 67. 3 mph 68. $-\frac{1}{28}$

Chapter 4 cumulative test

1. 25 2. -7 3. $\frac{4}{15}$ 4. 5 5. $\frac{13}{30}$ 6. $7x+11$
7. $-24a^6b^5$ 8. $x^3+2x^2-3x+20$ 9. 2 10. $\frac{x^2+6x-16}{x^2+3x}$
11. $\frac{-10y-13}{24}$ 12. $10x^2+39x-27$ 13. $16x^2-40x+25$
14. $9y^2-25$ 15. $2x^2+9x+27+\frac{80}{x-3}$
16. $\frac{6a^2+11a-10}{4a^2+23a-35}$ 17. $\frac{-2x+19}{(2x-3)(x+1)(2x+1)}$
18. $\{y|y \in R, y \neq \frac{3}{2}\}$ 19. $\{x|x \in R, x \neq \frac{1}{2}, -\frac{1}{2}\}$
20. $\{x|x \in R, x \neq -5, 5\}$ 21. $\{14\}$ 22. $\left\{-\frac{11}{24}\right\}$ 23. $\left\{-\frac{1}{8}\right\}$
24. $-\frac{13}{14}$ 25. $\frac{4x^2-3x-10}{5x^2+2x-16}$ 26. $\{x|x \leq -1\}$
27. $\{y|y < \frac{17}{2}\}$ 28. $\{z|z > -5\}$ 29. $\{x|x \geq \frac{97}{23}\}$
30. $-\frac{3a}{2b^2}$ 31. $\frac{p+4}{p+3}$ 32. $-\frac{3}{2}$ 33. $\left\{-\frac{1}{9}\right\}$ 34. $\frac{1}{2}$ or 6
35. $P(-2) = 27$

Chapter 5

Exercise 5-1

Answers to odd-numbered problems

1. 4.243 3. -5.745 5. $\sqrt[3]{9}$ 7. \sqrt{x} 9. $\sqrt[5]{b^4}$ 11. 4
13. 16 15. 27 17. 64 19. $\frac{1}{2}$ 21. $\frac{1}{2}$ 23. $\frac{1}{9}$ 25. $\frac{-1}{8}$
27. $\frac{1}{\sqrt[4]{x^3}}$ 29. $a^{4/7}$ 31. $x^{1/5}$ 33. -8 35. $|-4| = 4$
37. $|2x-y|$ 39. 256 41. $E = \frac{T}{\sqrt{(x^2+r^2)^3}}$
43. 24 miles per hour

Available for Fall 07 StudyBreaks
www.freeloadpress.com



Free Tunes.
For Students. By Students.

Student Musicians: Share your music. Make some change.

Learn more: www.freeloadpress.com/studybreakmusic

Contents

20 point learning system	xiii
Preface	xix
Study tips	xxv

Chapter 1 ■ Basic Concepts and Properties



1-1	Sets and real numbers	1
1-2	Operations with real numbers	12
1-3	Properties of real numbers	20
1-4	Order of operations	27
1-5	Terminology and evaluation	32
1-6	Sums and differences of polynomials	40
	Chapter 1 lead-in problem	46
	Chapter 1 summary	46
	Chapter 1 error analysis	47
	Chapter 1 critical thinking	47
	Chapter 1 review	47
	Chapter 1 test	49

Chapter 2 ■ First-Degree Equations and Inequalities



2-1	Solving equations	50
2-2	Formulas and literal equations	59
2-3	Word problems	63
2-4	Equations involving absolute value	72
2-5	Linear inequalities	77
2-6	Inequalities involving absolute value	86
	Chapter 2 lead-in problem	93
	Chapter 2 summary	93
	Chapter 2 error analysis	94
	Chapter 2 critical thinking	95
	Chapter 2 review	95
	Chapter 2 cumulative test	96

Chapter 3 ■ Exponents and Polynomials



3-1	Properties of exponents	97
3-2	Products of polynomials	103
3-3	Further properties of exponents	111
3-4	Common factors and factoring by grouping	121
3-5	Factoring trinomials of the form $x^2 + bx + c$ and perfect square trinomials	126
3-6	Factoring trinomials of the form $ax^2 + bx + c$	133
3-7	Other methods of factoring	141
3-8	Factoring: A general strategy	147
	Chapter 3 lead-in problem	150
	Chapter 3 summary	151
	Chapter 3 error analysis	151
	Chapter 3 critical thinking	152
	Chapter 3 review	152
	Chapter 3 cumulative test	153

Chapter 4 ■ Rational Expressions



4-1	Fundamental principle of rational expressions	154
4-2	Multiplication and division of rational expressions	160
4-3	Addition and subtraction of rational expressions	166
4-4	Complex rational expressions	176
4-5	Quotients of polynomials	183
4-6	Synthetic division, the remainder theorem, and the factor theorem	188
4-7	Equations containing rational expressions	198
4-8	Problem solving with rational equations	203
	Chapter 4 lead-in problem	209
	Chapter 4 summary	210
	Chapter 4 error analysis	211
	Chapter 4 critical thinking	211
	Chapter 4 review	212
	Chapter 4 cumulative test	214

Chapter 5 ■ Exponents, Roots, and Radicals



5-1	Roots and rational exponents	215
5-2	Operations with rational exponents	223
5-3	Simplifying radicals—I	226
5-4	Simplifying radicals—II	232
5-5	Sums and differences of radicals	237
5-6	Further operations with radicals	242
5-7	Complex numbers	246
	Chapter 5 lead-in problem	254
	Chapter 5 summary	254
	Chapter 5 error analysis	254
	Chapter 5 critical thinking	255
	Chapter 5 review	255
	Chapter 5 cumulative test	256

Chapter 6 ■ Quadratic Equations and Inequalities



6-1	Solution by factoring and extracting roots	258
6-2	Solution by completing the square	266
6-3	Solution by quadratic formula	271
6-4	Applications of quadratic equations	278
6-5	Equations involving radicals	285
6-6	Equations that are quadratic in form	289
6-7	Quadratic and rational inequalities	293
	Chapter 6 lead-in problem	300
	Chapter 6 summary	301
	Chapter 6 error analysis	301
	Chapter 6 critical thinking	302
	Chapter 6 review	302
	Chapter 6 cumulative test	304

Chapter 7 ■ Linear Equations and Inequalities in Two Variables



7-1	The rectangular coordinate system	305
7-2	The distance formula and the slope of a line	313
7-3	Finding the equation of a line	327
7-4	Graphs of linear inequalities	337
	Chapter 7 lead-in problem	343
	Chapter 7 summary	343
	Chapter 7 error analysis	344
	Chapter 7 critical thinking	345
	Chapter 7 review	345
	Chapter 7 cumulative test	346

Chapter 8 ■ Systems of Linear Equations



8-1	Systems of linear equations in two variables	348
8-2	Applied problems using systems of linear equations	358
8-3	Systems of linear equations in three variables	367
8-4	Determinants	375
8-5	Solutions of systems of linear equations by determinants	380
8-6	Solving systems of linear equations by the augmented matrix method	388
	Chapter 8 lead-in problem	394
	Chapter 8 summary	395
	Chapter 8 error analysis	395
	Chapter 8 critical thinking	396
	Chapter 8 review	397
	Chapter 8 cumulative test	399

Chapter 9 ■ Conic Sections



9-1 The parabola	401
9-2 More about parabolas	411
9-3 The circle	414
9-4 The ellipse and the hyperbola	420
9-5 Systems of nonlinear equations	429
Chapter 9 lead-in problem	435
Chapter 9 summary	435
Chapter 9 error analysis	436
Chapter 9 critical thinking	436
Chapter 9 review	437
Chapter 9 cumulative test	438

Chapter 10 ■ Functions



10-1 Relations and functions	440
10-2 Functional notation	449
10-3 Special functions and their graphs	455
10-4 Inverse functions	460
10-5 Variation	468
Chapter 10 lead-in problem	475
Chapter 10 summary	475
Chapter 10 error analysis	475
Chapter 10 critical thinking	476
Chapter 10 review	476
Chapter 10 cumulative test	477

Chapter 11 ■ Exponential and Logarithmic Functions

11-1	The exponential function	479
11-2	The logarithm	485
11-3	Properties of logarithms	490
11-4	The common logarithms	496
11-5	Logarithms to the base e	500
11-6	Exponential equations	505
	Chapter 11 lead-in problem	507
	Chapter 11 summary	507
	Chapter 11 error analysis	508
	Chapter 11 critical thinking	509
	Chapter 11 review	509
	Chapter 11 cumulative test	511

Chapter 12 ■ Sequences and Series

12-1	Sequences	513
12-2	Series	518
12-3	Arithmetic sequences	523
12-4	Geometric sequences and series	529
12-5	Infinite geometric series	536
12-6	The binomial expansion	541
	Chapter 12 lead-in problem	546
	Chapter 12 summary	546
	Chapter 12 error analysis	547
	Chapter 12 critical thinking	547
	Chapter 12 review	548
	Final examination	550

Appendix	Answers and solutions	553
Index		633

Index

A

Abscissa of a point, 307
 Absolute value, 9–10
 equation, 72–75
 inequalities, 86–90, 340–41
 Addition of complex numbers, 249
 Addition of fractions, 166
 Addition of rational expressions, 166–68, 171
 Addition property of equality, 23, 51
 Addition property of inequality, 79
 Additive inverse property, 22
 Algebraic expression, 32
 term of, 32
 Algebraic notation, 36
 Antilogarithms, 497
 Approximately equal to, 8, 217
 Arithmetic sequence, 523–24
 common difference of, 523
 general term of, 523–24
 sum of the terms of, 525
 Associative property of addition, 22
 Associative property of multiplication, 22
 Asymptotes, 423–24, 481
 Augmented matrix, 388
 Axes, x and y , 306
 Axiom, 20
 Axis of symmetry, 402

B

Base, 15, 97
 like, 98
 Binomial, 33
 expansion of, 541–44
 square of a , 105–6
 Braces, 1, 14
 Brackets, 14

C

Cantor, Georg, 1
 Circle
 center of, 415
 definition of, 414
 equation of a , 415–16
 general form of the equation of a , 416
 radius of a , 415
 standard form of the equation of a , 415
 Clearing fractions, 54

Closure property
 of addition, 22
 of multiplication, 22
 Coefficient, 32
 numerical, 32
 Combining like terms, 41
 Common difference, 523
 Common factors, 121–24
 Common logarithm, 496–97
 Common ratio, 530
 Commutative property
 of addition, 22
 of multiplication, 22
 Completely factored form, 121–23
 Completing the square, 266–67
 Complex conjugates, 250
 Complex numbers, 248
 addition of, 249
 definition of, 248
 division of, 251
 multiplication of, 250
 operations with, 248–51
 standard form of, 248
 subtraction of, 249
 Complex rational expressions, 176
 primary denominator of, 176
 primary numerator of, 176
 secondary denominators of, 176
 simplifying a, 176–79
 Components, of ordered pairs, 306
 Composite number, 121
 Composition of functions, 451
 Compound inequality, 78
 Conditional equation, 50
 Conic sections, 400
 Conjugate factors, 243
 complex, 250
 Consistent and independent system of equations, 350
 Constant function, 456
 Constant of variation, 468
 Contradiction, 55
 Coordinate(s), 7
 of a point, 307
 Cramer's Rule, 381–84
 Critical number, 293
 Cubes
 difference of two, 143–44
 sum of two, 144–45

D

Decay formulas, 502
 Decrease, 8
 Degree, 33
 Dependent system of equations, 350
 Dependent variable, 441
 Determinant, 375
 of a matrix, 375
 minor of, 376
 3×3 , 376
 2×2 , 375
 Difference of two cubes, 143–44
 Difference of two squares, 107, 141–42
 Direct variation, 468
 Discriminant, 274–75
 Disjoint sets, 4
 Distance formula, 315
 Distributive property, 22, 103
 Division, 16
 of complex numbers, 251
 definition of, 16
 involving zero, 17
 of a polynomial by a monomial, 183
 of a polynomial by a polynomial, 184
 of rational expressions, 162
 of rational numbers, 162
 Division property of rational expressions, 162
 Domain, 5
 of a function, 444–45
 of a rational expression, 155
 of a relation, 441
 Double-negative property, 24

E

Elementary row operations, 388
 Element of a set, 1
 Elimination, solution by, 350–53
 Ellipse
 definition of, 420
 equation of an, 421
 Empty set, 3
 Equality, 20
 Equality properties of real numbers, 21
 addition property, 23, 51
 multiplication property, 24, 52
 reflexive property, 21
 substitution property, 21, 34
 symmetric property, 21
 transitive property, 21

Equation, 50
 absolute value, 72–75
 of a circle, 415, 416
 conditional, 50
 of an ellipse, 421
 equivalent, 51
 exponential, 482, 505
 first-degree condition, 51
 graph of an, 308, 317
 of a hyperbola, 423
 of a line, 328
 linear, 51
 literal, 59
 logarithmic, 487
 nonlinear, 429
 of a parabola, 403, 413
 of quadratic form, 289
 root of an, 50
 solution of an, 50
 solving an, 53
 x-intercept of, 309
 y-intercept of, 309
 Equivalent equations, 51
 Evaluation, 34
 Expanded form, 15
 Exponential decay, 481, 502–3
 Exponential equation, 482, 505
 property of, 482
 Exponential form, 15, 97
 Exponential function, 479–81
 definition of, 479
 graph of, 480–81
 Exponential growth, 481, 502–3
 Exponential notation, 15, 97
 Exponents, 15
 definition, 97
 fraction to a power, 115–16
 group of factors to a power, 100
 negative, 112–13
 power of a power, 99
 product property, 98–99
 quotient property of, 111–12
 rational, 218–21, 223–25
 zero, 114
 Expression, algebraic, 32
 Extended distributive property, 103
 Extracting roots, 261
 Extraneous solutions, 199, 255

F

Factorial notation, 542
 Factoring, 121
 difference of two cubes, 143–44
 difference of two squares, 141–42
 four-term polynomials, 124–25
 a general strategy, 147–49
 greatest common factor, 121–22
 by grouping, 124–25
 by inspection, 136–40
 perfect-square trinomials, 130
 sum of two cubes, 144–45
 trinomials, 126–40

Factors, 14
 common, 121–24
 completely factored form, 121, 123
 conjugate, 242–43
 greatest common, 121–22
 prime factored form, 121
 Factor theorem, 192
 Finite, 4
 First component of an ordered pair, 306
 First-degree conditional equation, 51
 Foil, 104
 Formula, 59
 Function, 443
 composition of, 451
 constant, 456
 definition of, 443
 domain of, 443–45
 exponential, 479–81
 inverse, 460–63
 linear, 455
 logarithmic, 485
 notation, 449
 one-to-one, 461–62
 polynomial, 457
 quadratic, 456
 range of, 443
 square root, 458
 Fundamental principle of rational expressions, 156

G

General term
 of an arithmetic sequence, 523–24
 of a geometric sequence, 530
 of a sequence, 514
 Geometric formulas, Inside front cover
 Geometric sequence, 529
 common ratio of, 530
 sum of the terms of, 532
 Geometry problems, 66
 Graph, 7
 of a circle, 416–18
 of an ellipse, 422, 423
 of an equation, 308–11
 of a hyperbola, 425
 of linear inequalities in two variables, 337–40
 of a parabola, 404–7, 411–13
 Greater than, 8
 or equal to, 9
 Greatest common factor, 121–22
 Grouping symbols, 14, 42
 removing, 42
 Growth formula, 502

H

Horizontal line, slope of a, 320
 Horizontal line test, 462
 Hyperbola, 422
 asymptotes of, 423–24
 definition of, 422
 equation of, 423
 graph of, 425

I

Identical equation, 50
 Identity, 50
 property of addition, 22
 property of multiplication, 22
 Imaginary numbers, 246–48
 Inconsistent system of equations, 350
 Increase, 8
 Independent variable, 441
 Indeterminate, 17
 Index of summation, 519
 Inequalities
 absolute value, 86–90, 340–41
 addition property of, 79
 compound, 78
 is greater than, 8, 83
 is greater than or equal to, 9, 83
 is less than, 8, 83
 is less than or equal to, 9, 83
 linear, 77
 multiplication property of, 79–80
 order of, 80
 rational, 296
 sense of, 80
 solution set, 77–79
 strict, 8
 weak, 8
 Inequality properties of real numbers, 21
 transitive property, 21
 trichotomy property, 21
 Infinite, 4
 Infinite series, 536
 geometric, 536–38
 Infinity, 79
 Integer, 5
 Interest, simple, 65, 69
 Interest problem, 65, 69
 Intersection of sets, 3
 Interval notation, 78–79
 Inverse of a function, 460–63
 Inverse variation, 470
 Irrational numbers, 6, 217

J

Joint variation, 471

L

Least common denominator, 54
 Least common multiple, 168
 Left member, 50
 Less than, 8
 or equal to, 9
 Like bases, 98
 Like radicals, 237
 Like terms, 41
 Line, slope of a, 316–20
 Linear equation, 51
 systems of, 348
 in two variables, 305
 Linear function, 455
 Linear inequality, 77, 337
 graphs of, 337–40
 in two variables, 337

Line segment, 313
 midpoint of a, 316
 Listing method for sets, 1
 Literal equation, 59
 solving a, 60
 Logarithm, 485
 common, 496–97
 definition of, 485
 graph of, 485–86
 natural, 500
 power property of, 492
 product property of, 490
 quotient property of, 491
 Logarithmic
 equations, 487
 function, 485
 function, graph of, 485–86
 properties of, 487, 490–93
 Lower limit of summation, 519
 Lowest terms, reducing to, 156

M

Mathematical statement, 50
 Matrix, 375
 augmented, 388
 columns of, 375
 elements of, 375
 rows of, 375
 square, 375
 Member of an equation, 50
 Member of a set, 1
 Midpoint of a line segment, 316
 Minor of a determinant, 376
 Mixture problems, 71
 Monomial, 33
 Multinomial, 33
 multiplication of, 103–4, 108
 Multiplication, 15
 of fractions, 160
 of multinomials, 103–4, 108
 of rational expressions, 160
 of real numbers, 15
 Multiplication property of equality, 24, 52
 Multiplication property of inequality, 79–80
 Multiplication property of rational expressions, 160
 Multiplicative inverse property, 22
 Multiplicity, 193

N

Natural logarithms, 500
 Natural numbers, 4
 Negative exponents, 112–13
 Negative numbers, 5
 Negative reciprocal, 322
 n factorial, 542
 Nonlinear equations, systems of, 429–30
 n th power property, 255
 n th root, 215–17
 Null set, 3
 Number, 8
 Number line, 7
 Number problems, 64–65
 Numerical coefficient, 32

O

One-to-one
 function, 462
 Opposite of, 9
 Order, 8
 Ordered pairs of numbers, 306
 components of, 306
 Ordered triple of real numbers, 367
 Order of operations, 27–29
 Order relationship, 8, 80
 Ordinate of a point, 307
 Origin, 7, 306

P

Parabola, 401, 411
 definition of, 402
 equation of a, 402, 411
 vertex of a, 402
 Parallel lines, 321
 Parentheses, 14
 Partial sum of a series, 518
 Pascal's triangle, 541–42
 Perfect squares, 141
 trinomials, 130
 Perimeter, 66
 Perpendicular lines, 322
 Pi, 6, 32
 Plane, 400
 Point-slope form of a line, 328
 Polynomial, 33
 degree, 33
 division of, 183–85
 function, 457
 multiplication of, 103–8
 notation, 35
 sums and differences, 40–43
 Positive numbers, 4
 Primary
 denominator, 176
 numerator, 176
 Prime, relatively, 218
 Prime factor form, 121
 Prime numbers, 121
 Prime polynomial, 129
 Principal n th root, 216
 simplifying a, 227
 Problem solving, 29
 with linear equations, 64–66, 83
 with quadratic equations, 278–80
 with rational equations, 203–6
 with systems of linear equations, 358–60
 Product, 14
 Product property for radicals, 226
 Proof, 23
 Properties of a logarithm, 487, 490–93
 Properties of real numbers, 22
 Pythagorean Theorem, 208, 315

Q

Quadrants, 306
 Quadratic equation, 258
 applications of, 278–80
 in one variable, 258

solution by completing the square, 268–69
 solution by extracting roots, 261
 solution by factoring, 259
 solution by quadratic formula, 272–74
 standard form of, 258
 Quadratic formula, 272
 Quadratic function, 456
 Quadratic inequalities, 293–97
 critical numbers of, 293
 test number of, 294
 Quadratic-type equations, 289–91
 Quotient property of exponents, 112

R

Radical equations, 255
 solution set of, 255–57
 Radicals
 conjugate factors, 242
 differences of, 237
 index of a, 216
 like, 237
 multiplication of, 242
 product property, 226
 quotient property, 232
 simplest form, 235
 standard form of, 235
 sums of, 237
 Radicand, 216
 Range
 of a function, 444
 of a relation, 441
 Rational equations, 198
 Rational exponents, 218–21, 223–25
 Rational expression
 definition, 154
 domain of a, 155
 Rational inequality, 296–97
 Rationalizing the denominator, 232–34, 243–44
 Rational number, 6
 Real number, properties of, 22
 additive inverse property of, 22
 associative property of addition, 22
 associative property of multiplication, 22
 closure property of addition, 22
 closure property of multiplication, 22
 commutative property of addition, 22
 commutative property of multiplication, 22
 distributive property, 22
 identity property of addition, 22
 identity property of multiplication, 22
 multiplicative inverse property, 22
 Real number line, 7
 Real numbers, 6
 addition of, 12
 division of, 16
 multiplication of, 14–15
 subtraction of, 13
 Reciprocal, 22, 52, 162
 Rectangular coordinate system, 306
 Reducing to lowest terms, 156, 157
 Reflexive property of equality, 21
 Relation, 440
 domain of, 441
 range of, 441
 Relatively prime, 218
 Remainder theorem, 191

Replacement set, 5
 Right member, 50
 Root
 of an equation, 50
 n th, 215–17
 principal n th, 216
 Roster method for sets, 1
 r th term of a binomial expansion, 466

S

Scientific notation, 116–18
 Secondary denominator, 176
 Second component of an ordered pair, 306
 Sense of an inequality, 80
 Sequence, 513
 arithmetic, 523–24
 finite, 513
 infinite, 513
 general term of a, 514–15, 523–24, 530
 geometric, 522–30
 infinite, 513
 Series, 518
 arithmetic, 525
 geometric, 531
 infinite geometric, 536–38
 Set, 1
 disjoin, 4
 element of, 1
 empty, 3
 finite, 4
 infinite, 4
 intersection, 3
 member of, 1
 null, 3
 replacement, 5
 solution, 50
 union, 3
 Set-builder notation, 5
 Set of real numbers, 6
 Set symbolism, 1–4
 Sigma notation, 519
 index of, 519
 lower limit of, 519
 upper limit of, 519
 Sign, 12
 Sign array, of a determinant, 378
 Slope-intercept form, 329
 Slope of a line, 317–20
 definition of, 317
 horizontal line, 320
 vertical line, 320
 Solution, 50
 by completing the square, 268–69
 by elimination, 350–53
 by extracting the roots, 261–62
 by factoring, 259
 by quadratic formula, 272–73
 of an equation, 50
 of quadratic equations, 274
 of quadratic form equations, 290–91
 of quadratic inequalities, 293–94
 of radical equations, 255–57

 of rational equations, 198–99
 of rational inequalities, 293–94
 set, 50
 by substitution, 353, 354
 of systems by determinants, 380–84
 Special products, 105–7
 Square of a binomial, 106
 Square root function, 458
 Square root property, 261
 Squares, difference of two, 107, 141–42
 Standard form of a trinomial, 133
 Standard form of the equation of a line, 328
 Statement, mathematical, 50
 Strict inequality, 8, 303
 Subscripts, 35
 Subset, 2
 Substitution, property of, 21, 34
 Substitution, solution by, 166–68, 171, 353–54
 Subtraction, 13
 Subtraction, definition of, 13
 Subtraction of
 fractions, 166
 rational expressions, 166–67, 171
 real numbers, 13
 Summation notation, 519
 Sum of two cubes, 144–45
 Symbols
 absolute value, 8
 intersect, 3
 is an element of, 2
 is approximately equal to, 8, 217
 is a subset of, 2
 is greater than, 8
 is greater than or equal to, 9
 is less than, 8
 is less than or equal to, 9
 minus sign, 13
 multiplication dot, 14
 negative infinity, 79
 “not”—slash mark, 2
 null set or empty set, 3
 π , 6, 32
 plus sign, 13
 positive infinity, 79
 principal n th root, 216
 set of integers, 5
 set of irrational numbers, 6
 set of natural numbers, 4
 set of rational numbers, 6
 set of real numbers, 6
 set of whole numbers, 4
 union, 3
 Symmetric property of equality, 21
 Symmetry, 9
 axis of, 402
 Synthetic division, 188–91
 Systems of linear equations, 348
 applications, 358–60
 consistent and independent, 358
 dependent, 350
 graphs of, 350
 inconsistent, 350
 solution by augmented matrix, 388–92
 solution by determinants, 380–84
 solution by elimination, 350–53
 solution by substitution, 353–54
 three equations in three variables, 367
 Systems of nonlinear equations, 429

T

Term, 32
 Term, like, 41
 Test number, 293
 Theorem, 23
 Transitive property of equality, 21
 Transitive property of inequality, 21
 Trichotomy property, 21
 Trinomial, 33
 factoring a, 126–40
 standard form of, 133
 Triple, ordered, 367

U

Undefined, 17
 Union of sets, 3
 Unit distance, 7
 Upper limit of summation, 519

V

Variable, 5
 Variation, 468
 constant of, 468
 direct, 468–69
 inverse, 470–71
 joint, 471–72
 Vertex, of a parabola, 402
 Vertical line, slope of, 320
 Vertical line test, 445

W

Weak inequality, 9
 Whole numbers, 4

X

x -axis, 306
 x -intercept, 309, 403

Y

y -axis, 306
 y -intercept, 309, 404

Z

Zero
 division by, 17
 as an exponent, 114
 Zero factor property, 24
 Zero product property, 155